



# MMWR<sup>TM</sup>

## Morbidity and Mortality Weekly Report

Recommendations and Reports

October 25, 2002 / Vol. 51 / No. RR-16

### **Guideline for Hand Hygiene in Health-Care Settings**

**Recommendations of the Healthcare Infection Control Practices  
Advisory Committee and the HICPAC/SHEA/APIC/IDSA  
Hand Hygiene Task Force**

**INSIDE: Continuing Education Examination**

**CENTERS FOR DISEASE CONTROL AND PREVENTION**

**SAFER • HEALTHIER • PEOPLE<sup>TM</sup>**

The *MMWR* series of publications is published by the Epidemiology Program Office, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

### SUGGESTED CITATION

Centers for Disease Control and Prevention. Guideline for Hand Hygiene in Health-Care Settings: Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force. *MMWR* 2002;51(No. RR-16):[inclusive page numbers].

#### Centers for Disease Control and Prevention

Julie L. Gerberding, M.D., M.P.H.

*Director*

David W. Fleming, M.D.

*Deputy Director for Science and Public Health*

Dixie E. Snider, Jr., M.D., M.P.H.

*Associate Director for Science*

#### Epidemiology Program Office

Stephen B. Thacker, M.D., M.Sc.

*Director*

#### Office of Scientific and Health Communications

John W. Ward, M.D.

*Director*

*Editor, MMWR Series*

Suzanne M. Hewitt, M.P.A.

*Managing Editor*

Rachel J. Wilson

Douglas W. Weatherwax

*Project Editors*

Malbea A. Heilman

Beverly J. Holland

*Visual Information Specialists*

Quang M. Doan

Erica R. Shaver

*Information Technology Specialists*

## CONTENTS

### Part I. Review of the Scientific Data Regarding

|   |      |
|---|------|
| Hand Hygiene .....  | 1    |
| Historical Perspective .....  | 1    |
| Normal Bacterial Skin Flora .....   | 2    |
| Physiology of Normal Skin .....   | 2    |
| Definition of Terms .....   | 3    |
| Evidence of Transmission of Pathogens on Hands .....                                  | 4    |
| Models of Hand Transmission .....   | 5    |
| Relation of Hand Hygiene and Acquisition<br>of Health-Care–Associated Pathogens ..... | 5    |
| Methods Used To Evaluate the Efficacy<br>of Hand-Hygiene Products .....               | 6    |
| Review of Preparations Used for Hand Hygiene .....                                    | 8    |
| Activity of Antiseptic Agents Against<br>Spore-Forming Bacteria .....                 | 16   |
| Reduced Susceptibility of Bacteria to Antiseptics .....                               | 17   |
| Surgical Hand Antisepsis .....  | 17   |
| Relative Efficacy of Plain Soap, Antiseptic<br>Soap/Detergent, and Alcohols .....     | 18   |
| Irritant Contact Dermatitis Resulting from<br>Hand-Hygiene Measures .....             | 18   |
| Proposed Methods for Reducing Adverse<br>Effects of Agents .....                      | 19   |
| Factors To Consider When Selecting<br>Hand-Hygiene Products .....                     | 20   |
| Hand-Hygiene Practices Among HCWs .....   | 21   |
| Lessons Learned from Behavioral Theories .....  | 25   |
| Methods Used To Promote Improved Hand Hygiene .....                                   | 26   |
| Efficacy of Promotion and Impact of Improved<br>Hand Hygiene .....                    | 27   |
| Other Policies Related to Hand Hygiene .....  | 29   |
| Hand-Hygiene Research Agenda .....  | 30   |
| Web-Based Hand-Hygiene Resources .....  | 30   |
| Part II. Recommendations .....  | 31   |
| Categories .....  | 31   |
| Recommendations .....   | 32   |
| Part III. Performance Indicators .....  | 34   |
| References .....  | 34   |
| Appendix .....  | 45   |
| Continuing Education Activity .....   | CE-1 |

# Guideline for Hand Hygiene in Health-Care Settings

## Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force

Prepared by  
John M. Boyce, M.D.<sup>1</sup>  
Didier Pittet, M.D.<sup>2</sup>  
<sup>1</sup>Hospital of Saint Raphael  
New Haven, Connecticut  
<sup>2</sup>University of Geneva  
Geneva, Switzerland

### Summary

*The Guideline for Hand Hygiene in Health-Care Settings provides health-care workers (HCWs) with a review of data regarding handwashing and hand antisepsis in health-care settings. In addition, it provides specific recommendations to promote improved hand-hygiene practices and reduce transmission of pathogenic microorganisms to patients and personnel in health-care settings. This report reviews studies published since the 1985 CDC guideline (Garner JS, Favero MS. CDC guideline for handwashing and hospital environmental control, 1985. Infect Control 1986;7:231–43) and the 1995 APIC guideline (Larson EL, APIC Guidelines Committee. APIC guideline for handwashing and hand antisepsis in health care settings. Am J Infect Control 1995;23:251–69) were issued and provides an in-depth review of hand-hygiene practices of HCWs, levels of adherence of personnel to recommended handwashing practices, and factors adversely affecting adherence. New studies of the in vivo efficacy of alcohol-based hand rubs and the low incidence of dermatitis associated with their use are reviewed. Recent studies demonstrating the value of multidisciplinary hand-hygiene promotion programs and the potential role of alcohol-based hand rubs in improving hand-hygiene practices are summarized. Recommendations concerning related issues (e.g., the use of surgical hand antiseptics, hand lotions or creams, and wearing of artificial fingernails) are also included.*

## Part I. Review of the Scientific Data Regarding Hand Hygiene

### Historical Perspective

For generations, handwashing with soap and water has been considered a measure of personal hygiene (1). The concept of cleansing hands with an antiseptic agent probably emerged in the early 19<sup>th</sup> century. As early as 1822, a French pharmacist demonstrated that solutions containing chlorides of lime or soda could eradicate the foul odors associated with human corpses and that such solutions could be used as disinfectants and antiseptics (2). In a paper published in 1825, this pharmacist stated that physicians and other persons attending patients with contagious diseases would benefit from moistening their hands with a liquid chloride solution (2).

In 1846, Ignaz Semmelweis observed that women whose babies were delivered by students and physicians in the First Clinic at the General Hospital of Vienna consistently had a

higher mortality rate than those whose babies were delivered by midwives in the Second Clinic (3). He noted that physicians who went directly from the autopsy suite to the obstetrics ward had a disagreeable odor on their hands despite washing their hands with soap and water upon entering the obstetrics clinic. He postulated that the puerperal fever that affected so many parturient women was caused by “cadaverous particles” transmitted from the autopsy suite to the obstetrics ward via the hands of students and physicians. Perhaps because of the known deodorizing effect of chlorine compounds, as of May 1847, he insisted that students and physicians clean their hands with a chlorine solution between each patient in the clinic. The maternal mortality rate in the First Clinic subsequently dropped dramatically and remained low for years. This intervention by Semmelweis represents the first evidence indicating that cleansing heavily contaminated hands with an antiseptic agent between patient contacts may reduce health-care–associated transmission of contagious diseases more effectively than handwashing with plain soap and water.

In 1843, Oliver Wendell Holmes concluded independently that puerperal fever was spread by the hands of health personnel (1). Although he described measures that could be taken to limit its spread, his recommendations had little impact on

The material in this report originated in the National Center for Infectious Diseases, James M. Hughes, M.D., Director; and the Division of Healthcare Quality Promotion, Steve Solomon, M.D., Acting Director.

obstetric practices at the time. However, as a result of the seminal studies by Semmelweis and Holmes, handwashing gradually became accepted as one of the most important measures for preventing transmission of pathogens in health-care facilities.

In 1961, the U. S. Public Health Service produced a training film that demonstrated handwashing techniques recommended for use by health-care workers (HCWs) (4). At the time, recommendations directed that personnel wash their hands with soap and water for 1–2 minutes before and after patient contact. Rinsing hands with an antiseptic agent was believed to be less effective than handwashing and was recommended only in emergencies or in areas where sinks were unavailable.

In 1975 and 1985, formal written guidelines on handwashing practices in hospitals were published by CDC (5,6). These guidelines recommended handwashing with non-antimicrobial soap between the majority of patient contacts and washing with antimicrobial soap before and after performing invasive procedures or caring for patients at high risk. Use of waterless antiseptic agents (e.g., alcohol-based solutions) was recommended only in situations where sinks were not available.

In 1988 and 1995, guidelines for handwashing and hand antisepsis were published by the Association for Professionals in Infection Control (APIC) (7,8). Recommended indications for handwashing were similar to those listed in the CDC guidelines. The 1995 APIC guideline included more detailed discussion of alcohol-based hand rubs and supported their use in more clinical settings than had been recommended in earlier guidelines. In 1995 and 1996, the Healthcare Infection Control Practices Advisory Committee (HICPAC) recommended that either antimicrobial soap or a waterless antiseptic agent be used for cleaning hands upon leaving the rooms of patients with multidrug-resistant pathogens (e.g., vancomycin-resistant enterococci [VRE] and methicillin-resistant *Staphylococcus aureus* [MRSA]) (9,10). These guidelines also provided recommendations for handwashing and hand antisepsis in other clinical settings, including routine patient care. Although the APIC and HICPAC guidelines have been adopted by the majority of hospitals, adherence of HCWs to recommended handwashing practices has remained low (11,12).

Recent developments in the field have stimulated a review of the scientific data regarding hand hygiene and the development of new guidelines designed to improve hand-hygiene practices in health-care facilities. This literature review and accompanying recommendations have been prepared by a Hand Hygiene Task Force, comprising representatives from HICPAC, the Society for Healthcare Epidemiology of America (SHEA), APIC, and the Infectious Diseases Society of America (IDSA).

## Normal Bacterial Skin Flora

To understand the objectives of different approaches to hand cleansing, a knowledge of normal bacterial skin flora is essential. Normal human skin is colonized with bacteria; different areas of the body have varied total aerobic bacterial counts (e.g.,  $1 \times 10^6$  colony forming units (CFUs)/cm<sup>2</sup> on the scalp,  $5 \times 10^5$  CFUs/cm<sup>2</sup> in the axilla,  $4 \times 10^4$  CFUs/cm<sup>2</sup> on the abdomen, and  $1 \times 10^4$  CFUs/cm<sup>2</sup> on the forearm) (13). Total bacterial counts on the hands of medical personnel have ranged from  $3.9 \times 10^4$  to  $4.6 \times 10^6$  (14–17). In 1938, bacteria recovered from the hands were divided into two categories: transient and resident (14). Transient flora, which colonize the superficial layers of the skin, are more amenable to removal by routine handwashing. They are often acquired by HCWs during direct contact with patients or contact with contaminated environmental surfaces within close proximity of the patient. Transient flora are the organisms most frequently associated with health-care-associated infections. Resident flora, which are attached to deeper layers of the skin, are more resistant to removal. In addition, resident flora (e.g., coagulase-negative staphylococci and diphtheroids) are less likely to be associated with such infections. The hands of HCWs may become persistently colonized with pathogenic flora (e.g., *S. aureus*), gram-negative bacilli, or yeast. Investigators have documented that, although the number of transient and resident flora varies considerably from person to person, it is often relatively constant for any specific person (14,18).

## Physiology of Normal Skin

The primary function of the skin is to reduce water loss, provide protection against abrasive action and microorganisms, and act as a permeability barrier to the environment. The basic structure of skin includes, from outer- to innermost layer, the superficial region (i.e., the stratum corneum or horny layer, which is 10- to 20- $\mu$ m thick), the viable epidermis (50- to 100- $\mu$ m thick), the dermis (1- to 2-mm thick), and the hypodermis (1- to 2-mm thick). The barrier to percutaneous absorption lies within the stratum corneum, the thinnest and smallest compartment of the skin. The stratum corneum contains the corneocytes (or horny cells), which are flat, polyhedral-shaped nonnucleated cells, remnants of the terminally differentiated keratinocytes located in the viable epidermis. Corneocytes are composed primarily of insoluble bundled keratins surrounded by a cell envelope stabilized by cross-linked proteins and covalently bound lipid. Interconnecting the corneocytes of the stratum corneum are polar structures (e.g., corneodesmosomes), which contribute to stratum corneum cohesion.

The intercellular region of the stratum corneum is composed of lipid primarily generated from the exocytosis of lamellar bodies during the terminal differentiation of the keratinocytes. The intercellular lipid is required for a competent skin barrier and forms the only continuous domain. Directly under the stratum corneum is a stratified epidermis, which is composed primarily of 10–20 layers of keratinizing epithelial cells that are responsible for the synthesis of the stratum corneum. This layer also contains melanocytes involved in skin pigmentation; Langerhans cells, which are important for antigen presentation and immune responses; and Merkel cells, whose precise role in sensory reception has yet to be fully delineated. As keratinocytes undergo terminal differentiation, they begin to flatten out and assume the dimensions characteristic of the corneocytes (i.e., their diameter changes from 10–12  $\mu\text{m}$  to 20–30  $\mu\text{m}$ , and their volume increases by 10- to 20-fold). The viable epidermis does not contain a vascular network, and the keratinocytes obtain their nutrients from below by passive diffusion through the interstitial fluid.

The skin is a dynamic structure. Barrier function does not simply arise from the dying, degeneration, and compaction of the underlying epidermis. Rather, the processes of cornification and desquamation are intimately linked; synthesis of the stratum corneum occurs at the same rate as loss. Substantial evidence now confirms that the formation of the skin barrier is under homeostatic control, which is illustrated by the epidermal response to barrier perturbation by skin stripping or solvent extraction. Circumstantial evidence indicates that the rate of keratinocyte proliferation directly influences the integrity of the skin barrier. A general increase in the rate of proliferation results in a decrease in the time available for 1) uptake of nutrients (e.g., essential fatty acids), 2) protein and lipid synthesis, and 3) processing of the precursor molecules required for skin-barrier function. Whether chronic but quantitatively smaller increases in rate of epidermal proliferation also lead to changes in skin-barrier function remains unclear. Thus, the extent to which the decreased barrier function caused by irritants is caused by an increased epidermal proliferation also is unknown.

The current understanding of the formation of the stratum corneum has come from studies of the epidermal responses to perturbation of the skin barrier. Experimental manipulations that disrupt the skin barrier include 1) extraction of skin lipids with apolar solvents, 2) physical stripping of the stratum corneum using adhesive tape, and 3) chemically induced irritation. All of these experimental manipulations lead to a decreased skin barrier as determined by transepidermal water loss (TEWL). The most studied experimental system is the treatment of mouse skin with acetone. This experiment

results in a marked and immediate increase in TEWL, and therefore a decrease in skin-barrier function. Acetone treatment selectively removes glycerolipids and sterols from the skin, which indicates that these lipids are necessary, though perhaps not sufficient in themselves, for barrier function. Detergents act like acetone on the intercellular lipid domain. The return to normal barrier function is biphasic: 50%–60% of barrier recovery typically occurs within 6 hours, but complete normalization of barrier function requires 5–6 days.

## Definition of Terms

*Alcohol-based hand rub.* An alcohol-containing preparation designed for application to the hands for reducing the number of viable microorganisms on the hands. In the United States, such preparations usually contain 60%–95% ethanol or isopropanol.

*Antimicrobial soap.* Soap (i.e., detergent) containing an antiseptic agent.

*Antiseptic agent.* Antimicrobial substances that are applied to the skin to reduce the number of microbial flora. Examples include alcohols, chlorhexidine, chlorine, hexachlorophene, iodine, chloroxylenol (PCMX), quaternary ammonium compounds, and triclosan.

*Antiseptic handwash.* Washing hands with water and soap or other detergents containing an antiseptic agent.

*Antiseptic hand rub.* Applying an antiseptic hand-rub product to all surfaces of the hands to reduce the number of microorganisms present.

*Cumulative effect.* A progressive decrease in the numbers of microorganisms recovered after repeated applications of a test material.

*Decontaminate hands.* To Reduce bacterial counts on hands by performing antiseptic hand rub or antiseptic handwash.

*Detergent.* Detergents (i.e., surfactants) are compounds that possess a cleaning action. They are composed of both hydrophilic and lipophilic parts and can be divided into four groups: anionic, cationic, amphoteric, and nonionic detergents. Although products used for handwashing or antiseptic handwash in health-care settings represent various types of detergents, the term “soap” is used to refer to such detergents in this guideline.

*Hand antiseptics.* Refers to either antiseptic handwash or antiseptic hand rub.

*Hand hygiene.* A general term that applies to either handwashing, antiseptic handwash, antiseptic hand rub, or surgical hand antiseptics.

*Handwashing.* Washing hands with plain (i.e., non-antimicrobial) soap and water.



**Persistent activity.** Persistent activity is defined as the prolonged or extended antimicrobial activity that prevents or inhibits the proliferation or survival of microorganisms after application of the product. This activity may be demonstrated by sampling a site several minutes or hours after application and demonstrating bacterial antimicrobial effectiveness when compared with a baseline level. This property also has been referred to as “residual activity.” Both substantive and nonsubstantive active ingredients can show a persistent effect if they substantially lower the number of bacteria during the wash period.

**Plain soap.** Plain soap refers to detergents that do not contain antimicrobial agents or contain low concentrations of antimicrobial agents that are effective solely as preservatives.

**Substantivity.** Substantivity is an attribute of certain active ingredients that adhere to the stratum corneum (i.e., remain on the skin after rinsing or drying) to provide an inhibitory effect on the growth of bacteria remaining on the skin.

**Surgical hand antisepsis.** Antiseptic handwash or antiseptic hand rub performed preoperatively by surgical personnel to eliminate transient and reduce resident hand flora. Antiseptic detergent preparations often have persistent antimicrobial activity.

**Visibly soiled hands.** Hands showing visible dirt or visibly contaminated with proteinaceous material, blood, or other body fluids (e.g., fecal material or urine).

**Waterless antiseptic agent.** An antiseptic agent that does not require use of exogenous water. After applying such an agent, the hands are rubbed together until the agent has dried.

**Food and Drug Administration (FDA) product categories.** The 1994 FDA Tentative Final Monograph for Health-Care Antiseptic Drug Products divided products into three categories and defined them as follows (19):

- **Patient preoperative skin preparation.** A fast-acting, broad-spectrum, and persistent antiseptic-containing preparation that substantially reduces the number of microorganisms on intact skin.
- **Antiseptic handwash or HCW handwash.** An antiseptic-containing preparation designed for frequent use; it reduces the number of microorganisms on intact skin to an initial baseline level after adequate washing, rinsing, and drying; it is broad-spectrum, fast-acting, and if possible, persistent.
- **Surgical hand scrub.** An antiseptic-containing preparation that substantially reduces the number of microorganisms on intact skin; it is broad-spectrum, fast-acting, and persistent.

## Evidence of Transmission of Pathogens on Hands

Transmission of health-care-associated pathogens from one patient to another via the hands of HCWs requires the following sequence of events:

- Organisms present on the patient’s skin, or that have been shed onto inanimate objects in close proximity to the patient, must be transferred to the hands of HCWs.
- These organisms must then be capable of surviving for at least several minutes on the hands of personnel.
- Next, handwashing or hand antisepsis by the worker must be inadequate or omitted entirely, or the agent used for hand hygiene must be inappropriate.
- Finally, the contaminated hands of the caregiver must come in direct contact with another patient, or with an inanimate object that will come into direct contact with the patient.

Health-care-associated pathogens can be recovered not only from infected or draining wounds, but also from frequently colonized areas of normal, intact patient skin (20–31). The perineal or inguinal areas are usually most heavily colonized, but the axillae, trunk, and upper extremities (including the hands) also are frequently colonized (23,25,26,28,30–32). The number of organisms (e.g., *S. aureus*, *Proteus mirabilis*, *Klebsiella* spp., and *Acinetobacter* spp.) present on intact areas of the skin of certain patients can vary from 100 to 10<sup>6</sup>/cm<sup>2</sup> (25,29,31,33). Persons with diabetes, patients undergoing dialysis for chronic renal failure, and those with chronic dermatitis are likely to have areas of intact skin that are colonized with *S. aureus* (34–41). Because approximately 10<sup>6</sup> skin squames containing viable microorganisms are shed daily from normal skin (42), patient gowns, bed linen, bedside furniture, and other objects in the patient’s immediate environment can easily become contaminated with patient flora (30,43–46). Such contamination is particularly likely to be caused by staphylococci or enterococci, which are resistant to desiccation.

Data are limited regarding the types of patient-care activities that result in transmission of patient flora to the hands of personnel (26,45–51). In the past, attempts have been made to stratify patient-care activities into those most likely to cause hand contamination (52), but such stratification schemes were never validated by quantifying the level of bacterial contamination that occurred. Nurses can contaminate their hands with 100–1,000 CFUs of *Klebsiella* spp. during “clean” activities (e.g., lifting a patient; taking a patient’s pulse, blood pressure, or oral temperature; or touching a patient’s hand, shoulder, or groin) (48). Similarly, in another study, hands were cultured of nurses who touched the groins of patients heavily colonized with *P. mirabilis* (25); 10–600 CFUs/mL of this

organism were recovered from glove juice samples from the nurses' hands. Recently, other researchers studied contamination of HCWs' hands during activities that involved direct patient-contact wound care, intravascular catheter care, respiratory-tract care, and the handling of patient secretions (51). Agar fingertip impression plates were used to culture bacteria; the number of bacteria recovered from fingertips ranged from 0 to 300 CFUs. Data from this study indicated that direct patient contact and respiratory-tract care were most likely to contaminate the fingers of caregivers. Gram-negative bacilli accounted for 15% of isolates and *S. aureus* for 11%. Duration of patient-care activity was strongly associated with the intensity of bacterial contamination of HCWs' hands.

HCWs can contaminate their hands with gram-negative bacilli, *S. aureus*, enterococci, or *Clostridium difficile* by performing "clean procedures" or touching intact areas of the skin of hospitalized patients (26,45,46,53). Furthermore, personnel caring for infants with respiratory syncytial virus (RSV) infections have acquired RSV by performing certain activities (e.g., feeding infants, changing diapers, and playing with infants) (49). Personnel who had contact only with surfaces contaminated with the infants' secretions also acquired RSV by contaminating their hands with RSV and inoculating their oral or conjunctival mucosa. Other studies also have documented that HCWs may contaminate their hands (or gloves) merely by touching inanimate objects in patient rooms (46,53–56). None of the studies concerning hand contamination of hospital personnel were designed to determine if the contamination resulted in transmission of pathogens to susceptible patients.

Other studies have documented contamination of HCWs' hands with potential health-care-associated pathogens, but did not relate their findings to the specific type of preceding patient contact (15,17,57–62). For example, before glove use was common among HCWs, 15% of nurses working in an isolation unit carried a median of  $1 \times 10^4$  CFUs of *S. aureus* on their hands (61). Of nurses working in a general hospital, 29% had *S. aureus* on their hands (median count: 3,800 CFUs), whereas 78% of those working in a hospital for dermatology patients had the organism on their hands (median count:  $14.3 \times 10^6$  CFUs). Similarly, 17%–30% of nurses carried gram-negative bacilli on their hands (median counts: 3,400–38,000 CFUs). One study found that *S. aureus* could be recovered from the hands of 21% of intensive-care-unit personnel and that 21% of physician and 5% of nurse carriers had  $>1,000$  CFUs of the organism on their hands (59). Another study found lower levels of colonization on the hands of personnel working in a neurosurgery unit, with an average of 3 CFUs of *S. aureus* and 11 CFUs of gram-negative bacilli (16). Serial

cultures revealed that 100% of HCWs carried gram-negative bacilli at least once, and 64% carried *S. aureus* at least once.

## Models of Hand Transmission

Several investigators have studied transmission of infectious agents by using different experimental models. In one study, nurses were asked to touch the groins of patients heavily colonized with gram-negative bacilli for 15 seconds — as though they were taking a femoral pulse (25). Nurses then cleaned their hands by washing with plain soap and water or by using an alcohol hand rinse. After cleaning their hands, they touched a piece of urinary catheter material with their fingers, and the catheter segment was cultured. The study revealed that touching intact areas of moist skin of the patient transferred enough organisms to the nurses' hands to result in subsequent transmission to catheter material, despite handwashing with plain soap and water.

The transmission of organisms from artificially contaminated "donor" fabrics to clean "recipient" fabrics via hand contact also has been studied. Results indicated that the number of organisms transmitted was greater if the donor fabric or the hands were wet upon contact (63). Overall, only 0.06% of the organisms obtained from the contaminated donor fabric were transferred to recipient fabric via hand contact. *Staphylococcus saprophyticus*, *Pseudomonas aeruginosa*, and *Serratia* spp. were also transferred in greater numbers than was *Escherichia coli* from contaminated fabric to clean fabric after hand contact (64). Organisms are transferred to various types of surfaces in much larger numbers (i.e.,  $>10^4$ ) from wet hands than from hands that are thoroughly dried (65).

## Relation of Hand Hygiene and Acquisition of Health-Care-Associated Pathogens

Hand antisepsis reduces the incidence of health-care-associated infections (66,67). An intervention trial using historical controls demonstrated in 1847 that the mortality rate among mothers who delivered in the First Obstetrics Clinic at the General Hospital of Vienna was substantially lower when hospital staff cleaned their hands with an antiseptic agent than when they washed their hands with plain soap and water (3).

In the 1960s, a prospective, controlled trial sponsored by the National Institutes of Health and the Office of the Surgeon General demonstrated that infants cared for by nurses who did not wash their hands after handling an index infant colonized with *S. aureus* acquired the organism more often and more rapidly than did infants cared for by nurses who used hexachlorophene to clean their hands between infant

contacts (68). This trial provided evidence that, when compared with no handwashing, washing hands with an antiseptic agent between patient contacts reduces transmission of health-care-associated pathogens.

Trials have studied the effects of handwashing with plain soap and water versus some form of hand antiseptics on health-care-associated infection rates (69,70). Health-care-associated infection rates were lower when antiseptic handwashing was performed by personnel (69). In another study, antiseptic handwashing was associated with lower health-care-associated infection rates in certain intensive-care units, but not in others (70).

Health-care-associated infection rates were lower after antiseptic handwashing using a chlorhexidine-containing detergent compared with handwashing with plain soap or use of an alcohol-based hand rinse (71). However, because only a minimal amount of the alcohol rinse was used during periods when the combination regimen also was in use and because adherence to policies was higher when chlorhexidine was available, determining which factor (i.e., the hand-hygiene regimen or differences in adherence) accounted for the lower infection rates was difficult. Investigators have determined also that health-care-associated acquisition of MRSA was reduced when the antimicrobial soap used for hygienic handwashing was changed (72,73).

Increased handwashing frequency among hospital staff has been associated with decreased transmission of *Klebsiella* spp. among patients (48); these studies, however, did not quantify the level of handwashing among personnel. In a recent study, the acquisition of various health-care-associated pathogens was reduced when hand antiseptics was performed more frequently by hospital personnel (74); both this study and another (75) documented that the prevalence of health-care-associated infections decreased as adherence to recommended hand-hygiene measures improved.

Outbreak investigations have indicated an association between infections and understaffing or overcrowding; the association was consistently linked with poor adherence to hand hygiene. During an outbreak investigation of risk factors for central venous catheter-associated bloodstream infections (76), after adjustment for confounding factors, the patient-to-nurse ratio remained an independent risk factor for bloodstream infection, indicating that nursing staff reduction below a critical threshold may have contributed to this outbreak by jeopardizing adequate catheter care. The understaffing of nurses can facilitate the spread of MRSA in intensive-care settings (77) through relaxed attention to basic control measures (e.g., hand hygiene). In an outbreak of *Enterobacter cloacae* in a neonatal intensive-care unit (78), the daily number of

hospitalized children was above the maximum capacity of the unit, resulting in an available space per child below current recommendations. In parallel, the number of staff members on duty was substantially less than the number necessitated by the workload, which also resulted in relaxed attention to basic infection-control measures. Adherence to hand-hygiene practices before device contact was only 25% during the workload peak, but increased to 70% after the end of the understaffing and overcrowding period. Surveillance documented that being hospitalized during this period was associated with a fourfold increased risk of acquiring a health-care-associated infection. This study not only demonstrates the association between workload and infections, but it also highlights the intermediate cause of antimicrobial spread: poor adherence to hand-hygiene policies.

## Methods Used To Evaluate the Efficacy of Hand-Hygiene Products

### Current Methods

Investigators use different methods to study the in vivo efficacy of handwashing, antiseptic handwash, and surgical hand antiseptics protocols. Differences among the various studies include 1) whether hands are purposely contaminated with bacteria before use of test agents, 2) the method used to contaminate fingers or hands, 3) the volume of hand-hygiene product applied to the hands, 4) the time the product is in contact with the skin, 5) the method used to recover bacteria from the skin after the test solution has been used, and 6) the method of expressing the efficacy of the product (i.e., either percent reduction in bacteria recovered from the skin or log reduction of bacteria released from the skin). Despite these differences, the majority of studies can be placed into one of two major categories: studies focusing on products to remove transient flora and studies involving products that are used to remove resident flora from the hands. The majority of studies of products for removing transient flora from the hands of HCWs involve artificial contamination of the volunteer's skin with a defined inoculum of a test organism before the volunteer uses a plain soap, an antimicrobial soap, or a waterless antiseptic agent. In contrast, products tested for the preoperative cleansing of surgeons' hands (which must comply with surgical hand-antiseptics protocols) are tested for their ability to remove resident flora from without artificially contaminating the volunteers' hands.

In the United States, antiseptic handwash products intended for use by HCWs are regulated by FDA's Division of Over-the-Counter Drug Products (OTC). Requirements for in vitro and in vivo testing of HCW handwash products and surgical



hand scrubs are outlined in the FDA Tentative Final Monograph for Healthcare Antiseptic Drug Products (TFM) (19). Products intended for use as HCW handwashes are evaluated by using a standardized method (19). Tests are performed in accordance with use directions for the test material. Before baseline bacterial sampling and before each wash with the test material, 5 mL of a standardized suspension of *Serratia marcescens* are applied to the hands and then rubbed over the surfaces of the hands. A specified volume of the test material is dispensed into the hands and is spread over the hands and lower one third of the forearms. A small amount of tap water is added to the hands, and hands are completely lathered for a specified time, covering all surfaces of the hands and the lower third of the forearms. Volunteers then rinse hands and forearms under 40°C tap water for 30 seconds. Ten washes with the test formulation are required. After the first, third, seventh, and tenth washes, rubber gloves or polyethylene bags used for sampling are placed on the right and left hands, and 75 mL of sampling solution is added to each glove; gloves are secured above the wrist. All surfaces of the hand are massaged for 1 minute, and samples are obtained aseptically for quantitative culture. No neutralizer of the antimicrobial is routinely added to the sampling solution, but if dilution of the antimicrobial in the sampling fluid does not result in demonstrable neutralization, a neutralizer specific for the test formulation is added to the sampling solution. For waterless formulations, a similar procedure is used. TFM criteria for efficacy are as follows: a 2- $\log_{10}$  reduction of the indicator organism on each hand within 5 minutes after the first use, and a 3- $\log_{10}$  reduction of the indicator organism on each hand within 5 minutes after the tenth use (19).

Products intended for use as surgical hand scrubs have been evaluated also by using a standardized method (19). Volunteers clean under fingernails with a nail stick and clip their fingernails. All jewelry is removed from hands and arms. Hands and two thirds of forearms are rinsed with tap water (38°C–42°C) for 30 seconds, and then they are washed with a non-antimicrobial soap for 30 seconds and are rinsed for 30 seconds under tap water. Baseline microbial hand counts can then be determined. Next, a surgical scrub is performed with the test formulation using directions provided by the manufacturer. If no instructions are provided with the formulation, two 5-minute scrubs of hands and forearms followed by rinsing are performed. Reduction from baseline microbial hand counts is determined in a series of 11 scrubs conducted during 5 days. Hands are sampled at 1 minute, 3 hours, and 6 hours after the first scrubs on day 1, day 2, and day 5. After washing, volunteers wear rubber gloves; 75 mL of sampling solution are then added to one glove, and all surfaces of the hands are massaged

for 1 minute. Samples are then taken aseptically and cultured quantitatively. The other glove remains on the other hand for 6 hours and is sampled in the same manner. TFM requires that formulations reduce the number of bacteria 1  $\log_{10}$  on each hand within 1 minute of product application and that the bacterial cell count on each hand does not subsequently exceed baseline within 6 hours on day 1; the formulation must produce a 2- $\log_{10}$  reduction in microbial flora on each hand within 1 minute of product application by the end of the second day of enumeration and a 3- $\log_{10}$  reduction of microbial flora on each hand within 1 minute of product use by the end of the fifth day when compared with the established baseline (19).

The method most widely used in Europe to evaluate the efficacy of hand-hygiene agents is European Standard 1500–1997 (EN 1500—Chemical disinfectants and antiseptics. Hygienic hand-rub test method and requirements) (79). This method requires 12–15 test volunteers and an 18- to 24-hour growth of broth culture of *E. coli* K12. Hands are washed with a soft soap, dried, and then immersed halfway to the metacarpals in the broth culture for 5 seconds. Hands are removed from the broth culture, excess fluid is drained off, and hands are dried in the air for 3 minutes. Bacterial recovery for the initial value is obtained by kneading the fingertips of each hand separately for 60 seconds in 10 mL of tryptic soy broth (TSB) without neutralizers. The hands are removed from the broth and disinfected with 3 mL of the hand-rub agent for 30 seconds in a set design. The same operation is repeated with total disinfection time not exceeding 60 seconds. Both hands are rinsed in running water for 5 seconds and water is drained off. Fingertips of each hand are kneaded separately in 10 mL of TSB with added neutralizers. These broths are used to obtain the final value.  $\log_{10}$  dilutions of recovery medium are prepared and plated out. Within 3 hours, the same volunteers are tested with the reference disinfectant (60% 2-propanol [isopropanol]) and the test product. Colony counts are performed after 24 and 48 hours of incubation at 36°C. The average colony count of both left and right hand is used for evaluation. The log-reduction factor is calculated and compared with the initial and final values. The reduction factor of the test product should be superior or the same as the reference alcohol-based rub for acceptance. If a difference exists, then the results are analyzed statistically using the Wilcoxon test. Products that have log reductions substantially less than that observed with the reference alcohol-based hand rub (i.e., approximately 4  $\log_{10}$  reduction) are classified as not meeting the standard.

Because of different standards for efficacy, criteria cited in FDA TFM and the European EN 1500 document for establishing alcohol-based hand rubs vary (1, 19, 79). Alcohol-based

hand rubs that meet TFM criteria for efficacy may not necessarily meet the EN 1500 criteria for efficacy (80). In addition, scientific studies have not established the extent to which counts of bacteria or other microorganisms on the hands need to be reduced to minimize transmission of pathogens in health-care facilities (1,8); whether bacterial counts on the hands must be reduced by 1 log<sub>10</sub> (90% reduction), 2 log<sub>10</sub> (99%), 3 log<sub>10</sub> (99.9%), or 4 log<sub>10</sub> (99.99%) is unknown. Several other methods also have been used to measure the efficacy of antiseptic agents against various viral pathogens (81–83).

### Shortcomings of Traditional Methodologies

Accepted methods of evaluating hand-hygiene products intended for use by HCWs require that test volunteers wash their hands with a plain or antimicrobial soap for 30 seconds or 1 minute, despite the observation in the majority of studies that the average duration of handwashing by hospital personnel is <15 seconds (52,84–89). A limited number of investigators have used 15-second handwashing or hygienic hand-wash protocols (90–94). Therefore, almost no data exist regarding the efficacy of plain or antimicrobial soaps under conditions in which they are actually used by HCWs. Similarly, certain accepted methods for evaluating waterless antiseptic agents for use as antiseptic hand rubs require that 3 mL of alcohol be rubbed into the hands for 30 seconds, followed by a repeat application for the same duration. This type of protocol also does not reflect actual usage patterns among HCWs. Furthermore, volunteers used in evaluations of products are usually surrogates for HCWs, and their hand flora may not reflect flora found on the hands of personnel working in health-care settings. Further studies should be conducted among practicing HCWs using standardized protocols to obtain more realistic views of microbial colonization and risk of bacterial transfer and cross-transmission (51).

## Review of Preparations Used for Hand Hygiene

### Plain (Non-Antimicrobial) Soap

Soaps are detergent-based products that contain esterified fatty acids and sodium or potassium hydroxide. They are available in various forms including bar soap, tissue, leaflet, and liquid preparations. Their cleaning activity can be attributed to their detergent properties, which result in removal of dirt, soil, and various organic substances from the hands. Plain soaps have minimal, if any, antimicrobial activity. However, handwashing with plain soap can remove loosely adherent transient flora. For example, handwashing with plain soap and water for 15 seconds reduces bacterial counts on the skin by 0.6–1.1 log<sub>10</sub>, whereas washing for 30 seconds reduces counts

by 1.8–2.8 log<sub>10</sub> (1). However, in several studies, handwashing with plain soap failed to remove pathogens from the hands of hospital personnel (25,45). Handwashing with plain soap can result in paradoxical increases in bacterial counts on the skin (92,95–97). Non-antimicrobial soaps may be associated with considerable skin irritation and dryness (92,96,98), although adding emollients to soap preparations may reduce their propensity to cause irritation. Occasionally, plain soaps have become contaminated, which may lead to colonization of hands of personnel with gram-negative bacilli (99).

### Alcohols

The majority of alcohol-based hand antiseptics contain either isopropanol, ethanol, n-propanol, or a combination of two of these products. Although n-propanol has been used in alcohol-based hand rubs in parts of Europe for many years, it is not listed in TFM as an approved active agent for HCW handwashes or surgical hand-scrub preparations in the United States. The majority of studies of alcohols have evaluated individual alcohols in varying concentrations. Other studies have focused on combinations of two alcohols or alcohol solutions containing limited amounts of hexachlorophene, quaternary ammonium compounds, povidone-iodine, triclosan, or chlorhexidine gluconate (61,93,100–119).

The antimicrobial activity of alcohols can be attributed to their ability to denature proteins (120). Alcohol solutions containing 60%–95% alcohol are most effective, and higher concentrations are less potent (120–122) because proteins are not denatured easily in the absence of water (120). The alcohol content of solutions may be expressed as percent by weight (w/w), which is not affected by temperature or other variables, or as percent by volume (vol/vol), which can be affected by temperature, specific gravity, and reaction concentration (123). For example, 70% alcohol by weight is equivalent to 76.8% by volume if prepared at 15°C, or 80.5% if prepared at 25°C (123). Alcohol concentrations in antiseptic hand rubs are often expressed as percent by volume (19).

Alcohols have excellent in vitro germicidal activity against gram-positive and gram-negative vegetative bacteria, including multidrug-resistant pathogens (e.g., MRSA and VRE), *Mycobacterium tuberculosis*, and various fungi (120–122,124–129). Certain enveloped (lipophilic) viruses (e.g., herpes simplex virus, human immunodeficiency virus [HIV], influenza virus, respiratory syncytial virus, and vaccinia virus) are susceptible to alcohols when tested in vitro (120,130,131) (Table 1). Hepatitis B virus is an enveloped virus that is somewhat less susceptible but is killed by 60%–70% alcohol; hepatitis C virus also is likely killed by this percentage of alcohol (132). In a porcine tissue carrier model used to study antiseptic activity, 70% ethanol and 70% isopropanol were found to

**TABLE 1. Virucidal activity of antiseptic agents against enveloped viruses**

| Ref. no. | Test method                                     | Viruses               | Agent  | Results  |
|----------|---|-----------------------|--|--|
| (379)    | Suspension                                      | HIV                   | 19% EA   | LR = 2.0 in 5 minutes                                    |
| (380)    | Suspension                                      | HIV                   | 50% EA<br>35% IPA                                | LR > 3.5<br>LR > 3.7                                     |
| (381)    | Suspension                                      | HIV                   | 70% EA   | LR = 7.0 in 1 minute                                     |
| (382)    | Suspension                                      | HIV                   | 70% EA   | LR = 3.2B 5.5 in 30 seconds                              |
| (383)    | Suspension                                      | HIV                   | 70% IPA/0.5% CHG<br>4% CHG                       | LR = 6.0 in 15 seconds<br>LR = 6.0 in 15 seconds         |
| (384)    | Suspension                                      | HIV                   | Chloroxylenol<br>Benzalkonium chloride           | Inactivated in 1 minute<br>Inactivated in 1 minute       |
| (385)    | Suspension                                      | HIV                   | Povidone-iodine<br>Chlorhexidine                 | Inactivated<br>Inactivated                               |
| (386)    | Suspension                                      | HIV                   | Detergent/0.5%<br>PCMX                           | Inactivated in 30 seconds                                |
| (387)    | Suspension/dried plasma<br>chimpanzee challenge | HBV                   | 70% IPA  | LR = 6.0 in 10 minutes                                   |
| (388)    | Suspension/plasma<br>chimpanzee challenge       | HBV                   | 80% EA   | LR = 7.0 in 2 minutes                                    |
| (389)    | Suspension                                      | HSV                   | 95% EA<br>75% EA<br>95% IPA<br>70% EA + 0.5% CHG | LR > 5.0 in 1 minute<br>LR > 5.0<br>LR > 5.0<br>LR > 5.0 |
| (130)    | Suspension                                      | RSV                   | 35% IPA<br>4% CHG                                | LR > 4.3 in 1 minute<br>LR > 3.3                         |
| (141)    | Suspension                                      | Influenza<br>Vaccinia | 95% EA<br>95% EA                                 | Undetectable in 30 seconds<br>Undetectable in 30 seconds |
| (141)    | Hand test                                       | Influenza<br>Vaccinia | 95% EA<br>95% EA                                 | LR > 2.5<br>LR > 2.5                                     |

**Note:** HIV = human immunodeficiency virus, EA = ethanol, LR = Log<sub>10</sub> reduction, IPA = isopropanol, CHG = chlorhexidine gluconate, HBV = hepatitis B virus, RSV = respiratory syncytial virus, HSV = herpes simplex virus, HAV = hepatitis A virus, and PCMX = chloroxylenol.

reduce titers of an enveloped bacteriophage more effectively than an antimicrobial soap containing 4% chlorhexidine gluconate (133). Despite its effectiveness against these organisms, alcohols have very poor activity against bacterial spores, protozoan oocysts, and certain nonenveloped (nonlipophilic) viruses.

Numerous studies have documented the *in vivo* antimicrobial activity of alcohols. Alcohols effectively reduce bacterial counts on the hands (14, 121, 125, 134). Typically, log reductions of the release of test bacteria from artificially contaminated hands average 3.5 log<sub>10</sub> after a 30-second application and 4.0–5.0 log<sub>10</sub> after a 1-minute application (1). In 1994, the FDA TFM classified ethanol 60%–95% as a Category I agent (i.e., generally safe and effective for use in antiseptic handwash or HCW hand-wash products) (19). Although TFM placed isopropanol 70%–91.3% in category IIIIE (i.e., insufficient data to classify as effective), 60% isopropanol has subse-

quently been adopted in Europe as the reference standard against which alcohol-based hand-rub products are compared (79). Alcohols are rapidly germicidal when applied to the skin, but they have no appreciable persistent (i.e., residual) activity. However, regrowth of bacteria on the skin occurs slowly after use of alcohol-based hand antiseptics, presumably because of the sublethal effect alcohols have on some of the skin bacteria (135, 136). Addition of chlorhexidine, quaternary ammonium compounds, octenidine, or triclosan to alcohol-based solutions can result in persistent activity (1).

Alcohols, when used in concentrations present in alcohol-based hand rubs, also have *in vivo* activity against several nonenveloped viruses (Table 2). For example, 70% isopropanol and 70% ethanol are more effective than medicated soap or nonmedicated soap in reducing rotavirus titers on fingerpads (137, 138). A more recent study using the same test methods evaluated a commercially available product containing 60%

TABLE 2. Virucidal activity of antiseptic agents against nonenveloped viruses

| Ref. no. | Test method | Viruses                               | Antiseptic   | Result   |
|----------|-------------|---------------------------------------|--|--|
| (390)    | Suspension  | Rotavirus                             | 4% CHG<br>10% Povidone-Iodine<br>70% IPA/0.1% HCP  | LR < 3.0 in 1 minute<br>LR > 3.0<br>LR > 3.0   |
| (141)    | Hand test   | Adenovirus<br>Poliovirus<br>Coxsackie | 95% EA<br>95% EA<br>95% EA   | LR > 1.4<br>LR = 0.2–1.0<br>LR = 1.1–1.3   |
|          | Finger test | Adenovirus<br>Poliovirus<br>Coxsackie | 95% EA<br>95% EA<br>95% EA   | LR > 2.3<br>LR = 0.7–2.5<br>LR = 2.9   |
| (389)    | Suspension  | ECHO virus                            | 95% EA<br>75% EA<br>95% IPA<br>70% IPA + 0.5% CHG  | LR > 3.0 in 1 minute<br>LR ≤ 1.0<br>LR = 0<br>LR = 0   |
| (140)    | Finger pad  | HAV                                   | 70% EA<br>62% EA foam<br>plain soap<br>4% CHG<br>0.3% Triclosan  | 87.4% reduction<br>89.3% reduction<br>78.0% reduction<br>89.6% reduction<br>92.0% reduction                |
| (105)    | Finger tips | Bovine<br>Rotavirus                   | n-propanol + IPA<br>70% IPA<br>70% EA<br>2% triclosan<br>water (control)<br>7.5% povidone-iodine<br>plain soap<br>4% CHG | LR = 3.8 in 30 seconds<br>LR = 3.1<br>LR = 2.9<br>LR = 2.1<br>LR = 1.3<br>LR = 1.3<br>LR = 1.2<br>LR = 0.5 |
| (137)    | Finger pad  | Human<br>Rotavirus                    | 70% IPA<br>plain soap  | 98.9% decrease in 10 seconds<br>77.1%  |
| (138)    | Finger pad  | Human<br>Rotavirus                    | 70% IPA<br>2% CHG<br>plain soap  | 99.6% decrease in 10 seconds<br>80.3%<br>72.5%   |
| (81)     | Finger pad  | Rotavirus<br>Rhinovirus<br>Adenovirus | 60% EA gel<br>60% EA gel<br>60% EA gel   | LR > 3.0 in 10 seconds<br>LR > 3.0<br>LR > 3.0   |
| (139)    | Finger pad  | Poliovirus                            | 70% EA<br>70% IPA  | LR = 1.6 in 10 seconds<br>LR = 0.8   |
| (200)    | Finger tips | Poliovirus                            | Plain soap<br>80% EA   | LR = 2.1<br>LR = 0.4   |

**Note:** HIV = human immunodeficiency virus, EA = ethanol, LR = Log<sub>10</sub> reduction, IPA = isopropanol, CHG = chlorhexidine gluconate, HBV = hepatitis B virus, RSV = respiratory syncytial virus, HSV = herpes simplex virus, and HAV = hepatitis A virus.

ethanol and found that the product reduced the infectivity titers of three nonenveloped viruses (i.e., rotavirus, adenovirus, and rhinovirus) by >3 logs (81). Other nonenveloped viruses such as hepatitis A and enteroviruses (e.g., poliovirus) may require 70%–80% alcohol to be reliably inactivated (82,139). However, both 70% ethanol and a 62% ethanol foam product with emollients reduced hepatitis A virus titers on whole hands or fingertips more than nonmedicated soap; both were equally as effective as antimicrobial soap containing 4% chlorhexidine gluconate in reducing reduced viral counts on hands (140). In the same study, both 70% ethanol and the 62% ethanol foam product demonstrated greater virucidal activity against poliovirus than either non-antimicrobial

soap or a 4% chlorhexidine gluconate-containing soap (140). However, depending on the alcohol concentration, the amount of time that hands are exposed to the alcohol, and viral variant, alcohol may not be effective against hepatitis A and other nonlipophilic viruses. The inactivation of nonenveloped viruses is influenced by temperature, disinfectant-virus volume ratio, and protein load (141). Ethanol has greater activity against viruses than isopropanol. Further in vitro and in vivo studies of both alcohol-based formulations and antimicrobial soaps are warranted to establish the minimal level of virucidal activity that is required to interrupt direct contact transmission of viruses in health-care settings.



Alcohols are not appropriate for use when hands are visibly dirty or contaminated with proteinaceous materials. However, when relatively small amounts of proteinaceous material (e.g., blood) are present, ethanol and isopropanol may reduce viable bacterial counts on hands more than plain soap or antimicrobial soap (142).

Alcohol can prevent the transfer of health-care-associated pathogens (25,63,64). In one study, gram-negative bacilli were transferred from a colonized patient's skin to a piece of catheter material via the hands of nurses in only 17% of experiments after antiseptic hand rub with an alcohol-based hand rinse (25). In contrast, transfer of the organisms occurred in 92% of experiments after handwashing with plain soap and water. This experimental model indicates that when the hands of HCWs are heavily contaminated, an antiseptic hand rub using an alcohol-based rinse can prevent pathogen transmission more effectively than can handwashing with plain soap and water.

Alcohol-based products are more effective for standard handwashing or hand antisepsis by HCWs than soap or antimicrobial soaps (Table 3) (25,53,61,93,106–112,119,143–152). In all but two of the trials that compared alcohol-based solutions with antimicrobial soaps or detergents, alcohol reduced bacterial counts on hands more than washing hands with soaps or detergents containing hexachlorophene, povidone-iodine, 4% chlorhexidine, or triclosan. In studies exam-

ining antimicrobial-resistant organisms, alcohol-based products reduced the number of multidrug-resistant pathogens recovered from the hands of HCWs more effectively than did handwashing with soap and water (153–155).

Alcohols are effective for preoperative cleaning of the hands of surgical personnel (1,101,104,113–119,135,143,147,156–159) (Tables 4 and 5). In multiple studies, bacterial counts on the hands were determined immediately after using the product and again 1–3 hours later; the delayed testing was performed to determine if regrowth of bacteria on the hands is inhibited during operative procedures. Alcohol-based solutions were more effective than washing hands with plain soap in all studies, and they reduced bacterial counts on the hands more than antimicrobial soaps or detergents in the majority of experiments (101,104,113–119,135,143,147,157–159). In addition, the majority of alcohol-based preparations were more effective than povidone-iodine or chlorhexidine.

The efficacy of alcohol-based hand-hygiene products is affected by several factors, including the type of alcohol used, concentration of alcohol, contact time, volume of alcohol used, and whether the hands are wet when the alcohol is applied. Applying small volumes (i.e., 0.2–0.5 mL) of alcohol to the hands is not more effective than washing hands with plain soap and water (63,64). One study documented that 1 mL of alcohol was substantially less effective than 3 mL (91). The ideal volume of product to apply to the hands is not known

**TABLE 3. Studies comparing the relative efficacy (based on log<sub>10</sub> reductions achieved) of plain soap or antimicrobial soaps versus alcohol-based antiseptics in reducing counts of viable bacteria on hands**

| Ref. no. | Year | Skin contamination       | Assay method                | Time (sec) | Relative efficacy                                      |
|----------|------|--------------------------|-----------------------------|------------|--|
| (143)    | 1965 | Existing hand flora      | Finger-tip agar culture     | 60         | Plain soap < HCP < 50% EA foam                         |
| (119)    | 1975 | Existing hand flora      | Hand-rub broth culture      | —          | Plain soap < 95% EA                                    |
| (106)    | 1978 | Artificial contamination | Finger-tip broth culture    | 30         | Plain soap < 4% CHG < P-I < 70% EA = alc. CHG          |
| (144)    | 1978 | Artificial contamination | Finger-tip broth culture    | 30         | Plain soap < 4% CHG < 70% EA                           |
| (107)    | 1979 | Existing hand flora      | Hand-rub broth culture      | 120        | Plain soap < 0.5% aq. CHG < 70% EA < 4% CHG < alc.CHG  |
| (145)    | 1980 | Artificial contamination | Finger-tip broth culture    | 60–120     | 4% CHG < P-I < 60% IPA                                 |
| (53)     | 1980 | Artificial contamination | Finger-tip broth culture    | 15         | Plain soap < 3% HCP < P-I < 4% CHG < 70% EA            |
| (108)    | 1982 | Artificial contamination | Glove juice test            | 15         | P-I < alc. CHG   |
| (109)    | 1983 | Artificial contamination | Finger-tip broth culture    | 120        | 0.3–2% triclosan = 60% IPA = alc. CHG < alc. triclosan |
| (146)    | 1984 | Artificial contamination | Finger-tip agar culture     | 60         | Phenolic < 4% CHG < P-I < EA < IPA < n-P               |
| (147)    | 1985 | Existing hand flora      | Finger-tip agar culture     | 60         | Plain soap < 70% EA < 95% EA                           |
| (110)    | 1986 | Artificial contamination | Finger-tip broth culture    | 60         | Phenolic = P-I < alc. CHG < n-P                        |
| (93)     | 1986 | Existing hand flora      | Sterile-broth bag technique | 15         | Plain soap < IPA < 4% CHG = IPA-E = alc. CHG           |
| (61)     | 1988 | Artificial contamination | Finger-tip broth culture    | 30         | Plain soap < triclosan < P-I < IPA < alc. CHG < n-P    |
| (25)     | 1991 | Patient contact          | Glove-juice test            | 15         | Plain soap < IPA-E                                     |
| (148)    | 1991 | Existing hand flora      | Agar-plate/image analysis   | 30         | Plain soap < 1% triclosan < P-I < 4% CHG < IPA         |
| (111)    | 1992 | Artificial contamination | Finger-tip agar culture     | 60         | Plain soap < IPA < EA < alc. CHG                       |
| (149)    | 1992 | Artificial contamination | Finger-tip broth culture    | 60         | Plain soap < 60% n-P                                   |
| (112)    | 1994 | Existing hand flora      | Agar-plate/image analysis   | 30         | Plain soap < alc. CHG                                  |
| (150)    | 1999 | Existing hand flora      | Agar-plate culture          | N.S.       | Plain soap < commercial alcohol mixture                |
| (151)    | 1999 | Artificial contamination | Glove-juice test            | 20         | Plain soap < 0.6% PCMX < 65% EA                        |
| (152)    | 1999 | Artificial contamination | Finger-tip broth culture    | 30         | 4% CHG < plain soap < P-I < 70% EA                     |

**Note:** Existing hand flora = without artificially contaminating hands with bacteria, alc. CHG = alcoholic chlorhexidine gluconate, aq. CHG = aqueous chlorhexidine gluconate, 4% CHG = chlorhexidine gluconate detergent, EA = ethanol, HCP = hexachlorophene soap/detergent, IPA = isopropanol, IPA-E = isopropanol + emollients, n-P = n-propanol, PCMX = chloroxyleneol detergent, P-I = povidone-iodine detergent, and N.S. = not stated.

**TABLE 4. Studies comparing the relative efficacy of plain soap or antimicrobial soap versus alcohol-containing products in reducing counts of bacteria recovered from hands immediately after use of products for pre-operative cleansing of hands**

| Ref. no. | Year | Assay method               | Relative efficacy                             |
|----------|------|----------------------------|---|
| (143)    | 1965 | Finger-tip agar culture    | HCP < 50% EA foam + QAC                       |
| (157)    | 1969 | Finger-tip agar culture    | HCP < P-I < 50% EA foam + QAC                 |
| (101)    | 1973 | Finger-tip agar culture    | HCP soap < EA foam + 0.23% HCP                |
| (135)    | 1974 | Broth culture              | Plain soap < 0.5% CHG < 4% CHG < alc. CHG     |
| (119)    | 1975 | Hand-broth test            | Plain soap < 0.5% CHG < 4% CHG < alc. CHG     |
| (118)    | 1976 | Glove-juice test           | 0.5% CHG < 4% CHG < alc. CHG                  |
| (114)    | 1977 | Glove-juice test           | P-I < CHG < alc. CHG                          |
| (117)    | 1978 | Finger-tip agar culture    | P-I = 46% EA + 0.23% HCP                      |
| (113)    | 1979 | Broth culture of hands     | Plain soap < P-I < alc. CHG < alc. P-I        |
| (116)    | 1979 | Glove-juice test           | 70% IPA = alc. CHG                            |
| (147)    | 1985 | Finger-tip agar culture    | Plain soap < 70% - 90% EA                     |
| (115)    | 1990 | Glove-juice test, modified | Plain soap < triclosan < CHG < P-I < alc. CHG |
| (104)    | 1991 | Glove-juice test           | Plain soap < 2% triclosan < P-I < 70% IPA     |
| (158)    | 1998 | Finger-tip broth culture   | 70% IPA < 90% IPA = 60% n-P                   |
| (159)    | 1998 | Glove-juice test           | P-I < CHG < 70% EA                            |

**Note:** QAC = quaternary ammonium compound, alc. CHG = alcoholic chlorhexidine gluconate, CHG = chlorhexidine gluconate detergent, EA = ethanol, HCP = hexachlorophene detergent, IPA = isopropanol, and P-I = povidone-iodine detergent.

**TABLE 5. Efficacy of surgical hand-rub solutions in reducing the release of resident skin flora from clean hands**

| Study | Rub                                     | Concentration* (%) | Time (min)  | Mean log reduction                  |                  |                  |                  |                  |
|-------|---|--------------------|-------------|-------------------------------------|------------------|------------------|------------------|------------------|
|       |   |                    |             | Immediate                           | Sustained (3 hr) |                  |                  |                  |
| 1     | n-Propanol                              | 60                 | 5           | 2.9 <sup>†</sup>                    | 1.6 <sup>†</sup> |                  |                  |                  |
| 2     |   |                    | 5           | 2.7 <sup>†</sup>                    | NA               |                  |                  |                  |
| 3     |   |                    | 5           | 2.5 <sup>†</sup>                    | 1.8 <sup>†</sup> |                  |                  |                  |
| 4     |   |                    | 5           | 2.3 <sup>†</sup>                    | 1.6 <sup>†</sup> |                  |                  |                  |
| 5     |   |                    | 3           | 2.9 <sup>§</sup>                    | NA               |                  |                  |                  |
| 4     |   |                    | 3           | 2.0 <sup>†</sup>                    | 1.0 <sup>†</sup> |                  |                  |                  |
| 4     |   |                    | 1           | 1.1 <sup>†</sup>                    | 0.5 <sup>†</sup> |                  |                  |                  |
| 6     |   |                    | Isopropanol | 90                                  | 3                | 2.4 <sup>§</sup> | 1.4 <sup>§</sup> |                  |
| 6     |   |                    |             |                                     | 3                | 2.3 <sup>§</sup> | 1.2 <sup>§</sup> |                  |
| 7     |   |                    |             |                                     | 5                | 2.4 <sup>†</sup> | 2.1 <sup>†</sup> |                  |
| 4     | 5                                       | 2.1 <sup>†</sup>   |             |                                     | 1.0 <sup>†</sup> |                  |                  |                  |
| 6     | 3                                       | 2.0 <sup>§</sup>   |             |                                     | 0.7 <sup>§</sup> |                  |                  |                  |
| 5     | 3                                       | 1.7 <sup>c</sup>   |             |                                     | NA               |                  |                  |                  |
| 4     | 3                                       | 1.5 <sup>†</sup>   |             |                                     | 0.8 <sup>†</sup> |                  |                  |                  |
| 8     | 2                                       | 1.2                |             |                                     | 0.8              |                  |                  |                  |
| 4     | 1                                       | 0.7 <sup>†</sup>   | 0.2         |                                     |                  |                  |                  |                  |
| 9     | 1                                       | 0.8                | NA          |                                     |                  |                  |                  |                  |
| 10    | Isopropanol + chlorhexidine gluc. (w/v) | 60                 | 5           | 1.7                                 | 1.0              |                  |                  |                  |
| 7     |   |                    | 70 + 0.5    | 5                                   | 2.5 <sup>†</sup> | 2.7 <sup>†</sup> |                  |                  |
| 8     |   |                    |             | 2                                   | 1.0              | 1.5              |                  |                  |
| 11    |   |                    |             | Ethanol                             | 95               | 2.1              | NA               |                  |
| 5     |   |                    |             |                                     | 85               | 2.4 <sup>§</sup> | NA               |                  |
| 12    |   |                    |             |                                     | 80               | 1.5              | NA               |                  |
| 8     |   |                    |             |                                     | 70               | 1.0              | 0.6              |                  |
| 13    |   |                    |             | Ethanol + chlorhexidine gluc. (w/v) | 95 + 0.5         | 2                | 1.7              | NA               |
| 14    |   |                    |             |                                     | 77 + 0.5         | 5                | 2.0              | 1.5 <sup>¶</sup> |
| 8     |   |                    |             |                                     | 70 + 0.5         | 2                | 0.7              | 1.4              |
| 8     | Chlorhexidine gluc. (aq. Sol., w/v)     | 0.5                |             | 2                                   | 0.4              | 1.2              |                  |                  |
| 15    | Povidone-iodine (aq. Sol., w/v)         | 1.0                | 5           | 1.9 <sup>†</sup>                    | 0.8 <sup>†</sup> |                  |                  |                  |
| 16    | Peracetic acid (w/v)                    | 0.5                | 5           | 1.9                                 | NA               |                  |                  |                  |

**Note:** NA = not available.

**Source:** Rotter M. Hand washing and hand disinfection [Chapter 87]. In: Mayhall CG, ed. Hospital epidemiology and infection control. 2nd ed. Philadelphia, PA: Lippincott Williams & Wilkins, 1999. Table 5 is copyrighted by Lippincott Williams & Wilkins; it is reprinted here with their permission and permission from Manfred Rotler, M.D., Professor of Hygiene and Microbiology, Klinisches Institute für Hygiene der Universität Wien, Germany.

\* Volume/volume unless otherwise stated.

<sup>†</sup> Tested according to Deutsche Gesellschaft für Hygiene, and Mikrobiologic (DGHM)-German Society of Hygiene and Microbiology method.

<sup>§</sup> Tested according to European Standard prEN.

<sup>¶</sup> After 4 hours.

and may vary for different formulations. However, if hands feel dry after rubbing hands together for 10–15 seconds, an insufficient volume of product likely was applied. Because alcohol-impregnated towelettes contain a limited amount of alcohol, their effectiveness is comparable to that of soap and water (63,160,161).

Alcohol-based hand rubs intended for use in hospitals are available as low viscosity rinses, gels, and foams. Limited data are available regarding the relative efficacy of various formulations. One field trial demonstrated that an ethanol gel was slightly more effective than a comparable ethanol solution at reducing bacterial counts on the hands of HCWs (162). However, a more recent study indicated that rinses reduced bacterial counts on the hands more than the gels tested (80). Further studies are warranted to determine the relative efficacy of alcohol-based rinses and gels in reducing transmission of health-care-associated pathogens.

Frequent use of alcohol-based formulations for hand antisepsis can cause drying of the skin unless emollients, humectants, or other skin-conditioning agents are added to the formulations. The drying effect of alcohol can be reduced or eliminated by adding 1%–3% glycerol or other skin-conditioning agents (90,93,100,101,106,135,143,163,164). Moreover, in several recent prospective trials, alcohol-based rinses or gels containing emollients caused substantially less skin irritation and dryness than the soaps or antimicrobial detergents tested (96,98,165,166). These studies, which were conducted in clinical settings, used various subjective and objective methods for assessing skin irritation and dryness. Further studies are warranted to establish whether products with different formulations yield similar results.

Even well-tolerated alcohol hand rubs containing emollients may cause a transient stinging sensation at the site of any broken skin (e.g., cuts and abrasions). Alcohol-based hand-rub preparations with strong fragrances may be poorly tolerated by HCWs with respiratory allergies. Allergic contact dermatitis or contact urticaria syndrome caused by hypersensitivity to alcohol or to various additives present in certain alcohol hand rubs occurs only rarely (167,168).

Alcohols are flammable. Flash points of alcohol-based hand rubs range from 21°C to 24°C, depending on the type and concentration of alcohol present (169). As a result, alcohol-based hand rubs should be stored away from high temperatures or flames in accordance with National Fire Protection Agency recommendations. In Europe, where alcohol-based hand rubs have been used extensively for years, the incidence of fires associated with such products has been low (169). One recent U.S. report described a flash fire that occurred as a result of an unusual series of events, which included an HCW applying an alcohol gel to her hands, immediately removing a

polyester isolation gown, and then touching a metal door before the alcohol had evaporated (170). Removing the polyester gown created a substantial amount of static electricity that generated an audible static spark when the HCW touched the metal door, igniting the unevaporated alcohol on her hands (170). This incident emphasizes the need to rub hands together after application of alcohol-based products until all the alcohol has evaporated.

Because alcohols are volatile, containers should be designed to minimize evaporation. Contamination of alcohol-based solutions has seldom been reported. One report documented a cluster of pseudoinfections caused by contamination of ethyl alcohol by *Bacillus cereus* spores (171).

### Chlorhexidine

Chlorhexidine gluconate, a cationic bisbiguanide, was developed in England in the early 1950s and was introduced into the United States in the 1970s (8,172). Chlorhexidine base is only minimally soluble in water, but the digluconate form is water-soluble. The antimicrobial activity of chlorhexidine is likely attributable to attachment to, and subsequent disruption of, cytoplasmic membranes, resulting in precipitation of cellular contents (1,8). Chlorhexidine's immediate antimicrobial activity occurs more slowly than that of alcohols. Chlorhexidine has good activity against gram-positive bacteria, somewhat less activity against gram-negative bacteria and fungi, and only minimal activity against tubercle bacilli (1,8,172). Chlorhexidine is not sporicidal (1,172). It has in vitro activity against enveloped viruses (e.g., herpes simplex virus, HIV, cytomegalovirus, influenza, and RSV) but substantially less activity against nonenveloped viruses (e.g., rotavirus, adenovirus, and enteroviruses) (130,131,173). The antimicrobial activity of chlorhexidine is only minimally affected by the presence of organic material, including blood. Because chlorhexidine is a cationic molecule, its activity can be reduced by natural soaps, various inorganic anions, nonionic surfactants, and hand creams containing anionic emulsifying agents (8,172,174). Chlorhexidine gluconate has been incorporated into a number of hand-hygiene preparations. Aqueous or detergent formulations containing 0.5% or 0.75% chlorhexidine are more effective than plain soap, but they are less effective than antiseptic detergent preparations containing 4% chlorhexidine gluconate (135,175). Preparations with 2% chlorhexidine gluconate are slightly less effective than those containing 4% chlorhexidine (176).

Chlorhexidine has substantial residual activity (106,114–116,118,135,146,175). Addition of low concentrations (0.5%–1.0%) of chlorhexidine to alcohol-based preparations results in greater residual activity than alcohol alone (116,135). When used as recommended, chlorhexidine has a good safety

record (172). Minimal, if any, absorption of the compound occurs through the skin. Care must be taken to avoid contact with the eyes when using preparations with  $\geq 1\%$  chlorhexidine, because the agent can cause conjunctivitis and severe corneal damage. Ototoxicity precludes its use in surgery involving the inner or middle ear. Direct contact with brain tissue and the meninges should be avoided. The frequency of skin irritation is concentration-dependent, with products containing 4% most likely to cause dermatitis when used frequently for antiseptic handwashing (177); allergic reactions to chlorhexidine gluconate are uncommon (118,172). Occasional outbreaks of nosocomial infections have been traced to contaminated solutions of chlorhexidine (178–181).

### Chloroxylenol

Chloroxylenol, also known as parachlorometaxylenol (PCMX), is a halogen-substituted phenolic compound that has been used as a preservative in cosmetics and other products and as an active agent in antimicrobial soaps. It was developed in Europe in the late 1920s and has been used in the United States since the 1950s (182).

The antimicrobial activity of PCMX likely is attributable to inactivation of bacterial enzymes and alteration of cell walls (1). It has good in vitro activity against gram-positive organisms and fair activity against gram-negative bacteria, mycobacteria, and certain viruses (1,7,182). PCMX is less active against *P. aeruginosa*, but addition of ethylenediaminetetraacetic acid (EDTA) increases its activity against *Pseudomonas* spp. and other pathogens.

A limited number of articles focusing on the efficacy of PCMX-containing preparations intended for use by HCWs have been published in the last 25 years, and the results of studies have sometimes been contradictory. For example, in studies in which antiseptics were applied to abdominal skin, PCMX had the weakest immediate and residual activity of any of the agents studied (183). However, when 30-second handwashes were performed using 0.6% PCMX, 2% chlorhexidine gluconate, or 0.3% triclosan, the immediate effect of PCMX was similar to that of the other agents. When used 18 times per day for 5 consecutive days, PCMX had less cumulative activity than did chlorhexidine gluconate (184). When PCMX was used as a surgical scrub, one report indicated that 3% PCMX had immediate and residual activity comparable to 4% chlorhexidine gluconate (185), whereas two other studies demonstrated that the immediate and residual activity of PCMX was inferior to both chlorhexidine gluconate and povidone-iodine (176,186). The disparity between published studies may be associated with the various concentrations of PCMX included in the preparations evaluated and with other aspects of the formulations tested, including the

presence or absence of EDTA (7,182). PCMX is not as rapidly active as chlorhexidine gluconate or iodophors, and its residual activity is less pronounced than that observed with chlorhexidine gluconate (7,182). In 1994, FDA TFM tentatively classified PCMX as a Category III SE active agent (i.e., insufficient data are available to classify this agent as safe and effective) (19). Further evaluation of this agent by the FDA is ongoing.

The antimicrobial activity of PCMX is minimally affected by the presence of organic matter, but it is neutralized by non-ionic surfactants. PCMX, which is absorbed through the skin (7,182), is usually well-tolerated, and allergic reactions associated with its use are uncommon. PCMX is available in concentrations of 0.3%–3.75%. In-use contamination of a PCMX-containing preparation has been reported (187).

### Hexachlorophene

Hexachlorophene is a bisphenol composed of two phenolic groups and three chlorine moieties. In the 1950s and early 1960s, emulsions containing 3% hexachlorophene were widely used for hygienic handwashing, as surgical scrubs, and for routine bathing of infants in hospital nurseries. The antimicrobial activity of hexachlorophene results from its ability to inactivate essential enzyme systems in microorganisms. Hexachlorophene is bacteriostatic, with good activity against *S. aureus* and relatively weak activity against gram-negative bacteria, fungi, and mycobacteria (7).

Studies of hexachlorophene as a hygienic handwash and surgical scrub demonstrated only modest efficacy after a single handwash (53,143,188). Hexachlorophene has residual activity for several hours after use and gradually reduces bacterial counts on hands after multiple uses (i.e., it has a cumulative effect) (1,101,188,189). With repeated use of 3% hexachlorophene preparations, the drug is absorbed through the skin. Infants bathed with hexachlorophene and personnel regularly using a 3% hexachlorophene preparation for handwashing have blood levels of 0.1–0.6 ppm hexachlorophene (190). In the early 1970s, certain infants bathed with hexachlorophene developed neurotoxicity (vacuolar degeneration) (191). As a result, in 1972, the FDA warned that hexachlorophene should no longer be used routinely for bathing infants. However, after routine use of hexachlorophene for bathing infants in nurseries was discontinued, investigators noted that the incidence of health-care-associated *S. aureus* infections in hospital nurseries increased substantially (192,193). In several instances, the frequency of infections decreased when hexachlorophene bathing of infants was reinstated. However, current guidelines still recommend against the routine bathing of neonates with hexachlorophene because of its potential neurotoxic effects (194). The agent is classified by FDA TFM as not



generally recognized as safe and effective for use as an antiseptic handwash (19). Hexachlorophene should not be used to bathe patients with burns or extensive areas of susceptible, sensitive skin. Soaps containing 3% hexachlorophene are available by prescription only (7).

### Iodine and Iodophors

Iodine has been recognized as an effective antiseptic since the 1800s. However, because iodine often causes irritation and discoloring of skin, iodophors have largely replaced iodine as the active ingredient in antiseptics.

Iodine molecules rapidly penetrate the cell wall of microorganisms and inactivate cells by forming complexes with amino acids and unsaturated fatty acids, resulting in impaired protein synthesis and alteration of cell membranes (195). Iodophors are composed of elemental iodine, iodide or triiodide, and a polymer carrier (i.e., the complexing agent) of high molecular weight. The amount of molecular iodine present (so-called “free” iodine) determines the level of antimicrobial activity of iodophors. “Available” iodine refers to the total amount of iodine that can be titrated with sodium thiosulfate (196). Typical 10% povidone-iodine formulations contain 1% available iodine and yield free iodine concentrations of 1 ppm (196). Combining iodine with various polymers increases the solubility of iodine, promotes sustained release of iodine, and reduces skin irritation. The most common polymers incorporated into iodophors are polyvinyl pyrrolidone (i.e., povidone) and ethoxylated nonionic detergents (i.e., poloxamers) (195,196). The antimicrobial activity of iodophors also can be affected by pH, temperature, exposure time, concentration of total available iodine, and the amount and type of organic and inorganic compounds present (e.g., alcohols and detergents).

Iodine and iodophors have bactericidal activity against gram-positive, gram-negative, and certain spore-forming bacteria (e.g., clostridia and *Bacillus* spp.) and are active against mycobacteria, viruses, and fungi (8,195,197–200). However, in concentrations used in antiseptics, iodophors are not usually sporicidal (201). In vivo studies have demonstrated that iodophors reduce the number of viable organisms that are recovered from the hands of personnel (113,145,148,152,155). Povidone-iodine 5%–10% has been tentatively classified by FDA TFM as a Category I agent (i.e., a safe and effective agent for use as an antiseptic handwash and an HCW handwash) (19). The extent to which iodophors exhibit persistent antimicrobial activity after they have been washed off the skin is unclear. In one study, persistent activity was noted for 6 hours (176); however, several other studies demonstrated persistent activity for only 30–60 minutes after washing hands with an iodophor (61,117,202). In studies in which bacterial counts

were obtained after gloves were worn for 1–4 hours after washing, iodophors have demonstrated poor persistent activity (1,104,115,189,203–208). The in vivo antimicrobial activity of iodophors is substantially reduced in the presence of organic substances (e.g., blood or sputum) (8).

The majority of iodophor preparations used for hand hygiene contain 7.5%–10% povidone-iodine. Formulations with lower concentrations also have good antimicrobial activity because dilution can increase free iodine concentrations (209). However, as the amount of free iodine increases, the degree of skin irritation also may increase (209). Iodophors cause less skin irritation and fewer allergic reactions than iodine, but more irritant contact dermatitis than other antiseptics commonly used for hand hygiene (92). Occasionally, iodophor antiseptics have become contaminated with gram-negative bacilli as a result of poor manufacturing processes and have caused outbreaks or pseudo-outbreaks of infection (196).

### Quaternary Ammonium Compounds

Quaternary ammonium compounds are composed of a nitrogen atom linked directly to four alkyl groups, which may vary in their structure and complexity (210). Of this large group of compounds, alkyl benzalkonium chlorides are the most widely used as antiseptics. Other compounds that have been used as antiseptics include benzethonium chloride, cetrимide, and cetylpyridium chloride (1). The antimicrobial activity of these compounds was first studied in the early 1900s, and a quaternary ammonium compound for preoperative cleaning of surgeons' hands was used as early as 1935 (210). The antimicrobial activity of this group of compounds likely is attributable to adsorption to the cytoplasmic membrane, with subsequent leakage of low molecular weight cytoplasmic constituents (210).

Quaternary ammonium compounds are primarily bacteriostatic and fungistatic, although they are microbicidal against certain organisms at high concentrations (1); they are more active against gram-positive bacteria than against gram-negative bacilli. Quaternary ammonium compounds have relatively weak activity against mycobacteria and fungi and have greater activity against lipophilic viruses. Their antimicrobial activity is adversely affected by the presence of organic material, and they are not compatible with anionic detergents (1,210). In 1994, FDA TFM tentatively classified benzalkonium chloride and benzethonium chloride as Category IIISE active agents (i.e., insufficient data exists to classify them as safe and effective for use as an antiseptic handwash) (19). Further evaluation of these agents by FDA is in progress.

Quaternary ammonium compounds are usually well tolerated. However, because of weak activity against

gram-negative bacteria, benzalkonium chloride is prone to contamination by these organisms. Several outbreaks of infection or pseudoinfection have been traced to quaternary ammonium compounds contaminated with gram-negative bacilli (211–213). For this reason, in the United States, these compounds have been seldom used for hand antisepsis during the last 15–20 years. However, newer handwashing products containing benzalkonium chloride or benzethonium chloride have recently been introduced for use by HCWs. A recent study of surgical intensive-care unit personnel found that cleaning hands with antimicrobial wipes containing a quaternary ammonium compound was about as effective as using plain soap and water for handwashing; both were less effective than decontaminating hands with an alcohol-based hand rub (214). One laboratory-based study reported that an alcohol-free hand-rub product containing a quaternary ammonium compound was efficacious in reducing microbial counts on the hands of volunteers (215). Further studies of such products are needed to determine if newer formulations are effective in health-care settings.

### Triclosan

Triclosan (chemical name: 2,4,4'-trichloro-2'-hydroxydiphenyl ether) is a nonionic, colorless substance that was developed in the 1960s. It has been incorporated into soaps for use by HCWs and the public and into other consumer products. Concentrations of 0.2%–2% have antimicrobial activity. Triclosan enters bacterial cells and affects the cytoplasmic membrane and synthesis of RNA, fatty acids, and proteins (216). Recent studies indicate this agent's antibacterial activity is attributable to binding to the active site of enoyl-acyl carrier protein reductase (217,218).

Triclosan has a broad range of antimicrobial activity, but it is often bacteriostatic (1). Minimum inhibitory concentrations (MICs) range from 0.1 to 10 µg/mL, whereas minimum bactericidal concentrations are 25–500 µg/mL. Triclosan's activity against gram-positive organisms (including MRSA) is greater than against gram-negative bacilli, particularly *P. aeruginosa* (1,216). The agent possesses reasonable activity against mycobacterial and *Candida* spp., but it has limited activity against filamentous fungi. Triclosan (0.1%) reduces bacterial counts on hands by 2.8 log<sub>10</sub> after a 1-minute hygienic handwash (1). In several studies, log reductions have been lower after triclosan is used than when chlorhexidine, iodophors, or alcohol-based products are applied (1,61,149,184,219). In 1994, FDA TFM tentatively classified triclosan ≤1.0% as a Category IIISE active agent (i.e., insufficient data exist to classify this agent as safe and effective for use as an antiseptic handwash) (19). Further evaluation of this agent by the FDA is underway. Like chlorhexidine, triclosan has persistent activity on the skin. Its activity in

hand-care products is affected by pH, the presence of surfactants, emollients, or humectants and by the ionic nature of the particular formulation (1,216). Triclosan's activity is not substantially affected by organic matter, but it can be inhibited by sequestration of the agent in micelle structures formed by surfactants present in certain formulations. The majority of formulations containing <2% triclosan are well-tolerated and seldom cause allergic reactions. Certain reports indicate that providing hospital personnel with a triclosan-containing preparation for hand antisepsis has led to decreased MRSA infections (72,73). Triclosan's lack of potent activity against gram-negative bacilli has resulted in occasional reports of contamination (220).

### Other Agents

Approximately 150 years after puerperal-fever-related maternal mortality rates were demonstrated by Semmelweis to be reduced by use of a hypochlorite hand rinse, the efficacy of rubbing hands for 30 seconds with an aqueous hypochlorite solution was studied once again (221). The solution was demonstrated to be no more effective than distilled water. The regimen used by Semmelweis, which called for rubbing hands with a 4% [w/w] hypochlorite solution until the hands were slippery (approximately 5 minutes), has been revisited by other researchers (222). This more current study indicated that the regimen was 30 times more effective than a 1-minute rub using 60% isopropanol. However, because hypochlorite solutions are often irritating to the skin when used repeatedly and have a strong odor, they are seldom used for hand hygiene.

Certain other agents are being evaluated by FDA for use in health-care-related antiseptics (19). However, the efficacy of these agents has not been evaluated adequately for use in handwashing preparations intended for use by HCWs. Further evaluation of these agents is warranted. Products that use different concentrations of traditional antiseptics (e.g., low concentrations of iodophor) or contain novel compounds with antiseptic properties are likely to be introduced for use by HCWs. For example, preliminary studies have demonstrated that adding silver-containing polymers to an ethanol carrier (i.e., Surfacine<sup>®</sup>) results in a preparation that has persistent antimicrobial activity on animal and human skin (223). New compounds with good in vitro activity must be tested in vivo to determine their abilities to reduce transient and resident skin flora on the hands of HCWs.

### Activity of Antiseptic Agents Against Spore-Forming Bacteria

The widespread prevalence of health-care-associated diarrhea caused by *Clostridium difficile* and the recent occurrence

in the United States of human *Bacillus anthracis* infections associated with contaminated items sent through the postal system has raised concern regarding the activity of antiseptic agents against spore-forming bacteria. None of the agents (including alcohols, chlorhexidine, hexachlorophene, iodophors, PCMX, and triclosan) used in antiseptic handwash or antiseptic hand-rub preparations are reliably sporicidal against *Clostridium* spp. or *Bacillus* spp. (120,172,224,225). Washing hands with non-antimicrobial or antimicrobial soap and water may help to physically remove spores from the surface of contaminated hands. HCWs should be encouraged to wear gloves when caring for patients with *C. difficile*-associated diarrhea (226). After gloves are removed, hands should be washed with a non-antimicrobial or an antimicrobial soap and water or disinfected with an alcohol-based hand rub. During outbreaks of *C. difficile*-related infections, washing hands with a non-antimicrobial or antimicrobial soap and water after removing gloves is prudent. HCWs with suspected or documented exposure to *B. anthracis*-contaminated items also should be encouraged to wash their hands with a non-antimicrobial or antimicrobial soap and water.

### Reduced Susceptibility of Bacteria to Antiseptics

Reduced susceptibility of bacteria to antiseptic agents can either be an intrinsic characteristic of a species or can be an acquired trait (227). Several reports have described strains of bacteria that appear to have acquired reduced susceptibility (when defined by MICs established in vitro) to certain antiseptics (e.g., chlorhexidine, quaternary ammonium compounds, and triclosan) (227–230). However, because the antiseptic concentrations that are actually used by HCWs are often substantially higher than the MICs of strains with reduced antiseptic susceptibility, the clinical relevance of the in vitro findings is questionable. For example, certain strains of MRSA have chlorhexidine and quaternary ammonium compound MICs that are several-fold higher than methicillin-susceptible strains, and certain strains of *S. aureus* have elevated MICs to triclosan (227,228). However, such strains were readily inhibited by the concentrations of these antiseptics that are actually used by practicing HCWs (227,228). The description of a triclosan-resistant bacterial enzyme has raised the question of whether resistance to this agent may develop more readily than to other antiseptic agents (218). In addition, exposing *Pseudomonas* strains containing the MexAB-OprM efflux system to triclosan may select for mutants that are resistant to multiple antibiotics, including fluoroquinolones (230). Further studies are needed to determine whether reduced susceptibility to antiseptic agents is of epidemiologic

significance and whether resistance to antiseptics has any influence on the prevalence of antibiotic-resistant strains (227).

### Surgical Hand Antisepsis

Since the late 1800s, when Lister promoted the application of carbolic acid to the hands of surgeons before procedures, preoperative cleansing of hands and forearms with an antiseptic agent has been an accepted practice (231). Although no randomized, controlled trials have been conducted to indicate that surgical-site infection rates are substantially lower when preoperative scrubbing is performed with an antiseptic agent rather than a non-antimicrobial soap, certain other factors provide a strong rationale for this practice. Bacteria on the hands of surgeons can cause wound infections if introduced into the operative field during surgery (232); rapid multiplication of bacteria occurs under surgical gloves if hands are washed with a non-antimicrobial soap. However, bacterial growth is slowed after preoperative scrubbing with an antiseptic agent (14,233). Reducing resident skin flora on the hands of the surgical team for the duration of a procedure reduces the risk of bacteria being released into the surgical field if gloves become punctured or torn during surgery (1,156,169). Finally, at least one outbreak of surgical-site infections occurred when surgeons who normally used an antiseptic surgical scrub preparation began using a non-antimicrobial product (234).

Antiseptic preparations intended for use as surgical hand scrubs are evaluated for their ability to reduce the number of bacteria released from hands at different times, including 1) immediately after scrubbing, 2) after wearing surgical gloves for 6 hours (i.e., persistent activity), and 3) after multiple applications over 5 days (i.e., cumulative activity). Immediate and persistent activity are considered the most important in determining the efficacy of the product. U.S. guidelines recommend that agents used for surgical hand scrubs should substantially reduce microorganisms on intact skin, contain a nonirritating antimicrobial preparation, have broad-spectrum activity, and be fast-acting and persistent (19,235).

Studies have demonstrated that formulations containing 60%–95% alcohol alone or 50%–95% when combined with limited amounts of a quaternary ammonium compound, hexachlorophene, or chlorhexidine gluconate, lower bacterial counts on the skin immediately postscrub more effectively than do other agents (Table 4). The next most active agents (in order of decreasing activity) are chlorhexidine gluconate, iodophors, triclosan, and plain soap (104,119,186,188,203,204,206,208,236). Because studies of PCMX as a surgical scrub have yielded contradictory results, further studies are needed to establish how the efficacy of this compound compares with the other agents (176,185,186).

Although alcohols are not considered to have persistent antimicrobial activity, bacteria appear to reproduce slowly on the hands after a surgical scrub with alcohol, and bacterial counts on hands after wearing gloves for 1–3 hours seldom exceed baseline (i.e., prescrub) values (1). However, a recent study demonstrated that a formulation containing 61% ethanol alone did not achieve adequate persistent activity at 6 hours postscrub (237). Alcohol-based preparations containing 0.5% or 1% chlorhexidine gluconate have persistent activity that, in certain studies, has equaled or exceeded that of chlorhexidine gluconate-containing detergents (1,118,135,237).\*

Persistent antimicrobial activity of detergent-based surgical scrub formulations is greatest for those containing 2% or 4% chlorhexidine gluconate, followed by hexachlorophene, triclosan, and iodophors (1,102,113–115,159,189,203,204,206–208,236). Because hexachlorophene is absorbed into the blood after repeated use, it is seldom used as a surgical scrub.

Surgical staff have been traditionally required to scrub their hands for 10 minutes preoperatively, which frequently leads to skin damage. Several studies have demonstrated that scrubbing for 5 minutes reduces bacterial counts as effectively as a 10-minute scrub (117,238,239). In other studies, scrubbing for 2 or 3 minutes reduced bacterial counts to acceptable levels (156,205,207,240,241).

Studies have indicated that a two-stage surgical scrub using an antiseptic detergent, followed by application of an alcohol-containing preparation, is effective. For example, an initial 1- or 2-minute scrub with 4% chlorhexidine gluconate or povidone-iodine followed by application of an alcohol-based product has been as effective as a 5-minute scrub with an antiseptic detergent (114,242).

Surgical hand-antiseptic protocols have required personnel to scrub with a brush. But this practice can damage the skin of personnel and result in increased shedding of bacteria from the hands (95,243). Scrubbing with a disposable sponge or combination sponge-brush has reduced bacterial counts on the hands as effectively as scrubbing with a brush (244–246). However, several studies indicate that neither a brush nor a

sponge is necessary to reduce bacterial counts on the hands of surgical personnel to acceptable levels, especially when alcohol-based products are used (102,117,159,165,233,237,247,248). Several of these studies performed cultures immediately or at 45–60 minutes postscrub (102,117,233,247,248), whereas in other studies, cultures were obtained 3 and 6 hours postscrub (159,237). For example, a recent laboratory-based study using volunteers demonstrated that brushless application of a preparation containing 1% chlorhexidine gluconate plus 61% ethanol yielded lower bacterial counts on the hands of participants than using a sponge/brush to apply a 4% chlorhexidine-containing detergent preparation (237).

### **Relative Efficacy of Plain Soap, Antiseptic Soap/Detergent, and Alcohols**

Comparing studies related to the in vivo efficacy of plain soap, antimicrobial soaps, and alcohol-based hand rubs is problematic, because certain studies express efficacy as the percentage reduction in bacterial counts achieved, whereas others give log<sub>10</sub> reductions in counts achieved. However, summarizing the relative efficacy of agents tested in each study can provide an overview of the in vivo activity of various formulations intended for handwashing, hygienic handwash, antiseptic hand rub, or surgical hand antiseptics (Tables 2–4).

### **Irritant Contact Dermatitis Resulting from Hand-Hygiene Measures**

#### **Frequency and Pathophysiology of Irritant Contact Dermatitis**

In certain surveys, approximately 25% of nurses report symptoms or signs of dermatitis involving their hands, and as many as 85% give a history of having skin problems (249). Frequent and repeated use of hand-hygiene products, particularly soaps and other detergents, is a primary cause of chronic irritant contact dermatitis among HCWs (250). The potential of detergents to cause skin irritation can vary considerably and can be ameliorated by the addition of emollients and humectants. Irritation associated with antimicrobial soaps may be caused by the antimicrobial agent or by other ingredients of the formulation. Affected persons often complain of a feeling of dryness or burning; skin that feels “rough;” and erythema, scaling, or fissures. Detergents damage the skin by causing denaturation of stratum corneum proteins, changes in intercellular lipids (either depletion or reorganization of lipid moieties), decreased corneocyte cohesion, and decreased stratum corneum water-binding capacity (250,251). Damage

\* In a recent randomized clinical trial, surgical site infection rates were monitored among patients who were operated on by surgical personnel who cleaned their hands preoperatively either by performing a traditional 5-minute surgical hand scrub using 4% povidone-iodine or 4% antiseptic antimicrobial soap, or by washing their hands for 1 minute with a non-antimicrobial soap followed by a 5-minute hand-rubbing technique using an alcohol-based hand rinse containing 0.2% mectronium etilsulfate. The incidence of surgical site infections was virtually identical in the two groups of patients. (Source: Parienti JJ, Thibon P, Heller R, et al. for Members of the Antiseptic Chirurgicale des Mains Study Group. Hand-rubbing with an aqueous alcoholic solution vs traditional surgical hand-scrubbing and 30-day surgical site infection rates: a randomized equivalence study. JAMA 2002;288:722–7).



to the skin also changes skin flora, resulting in more frequent colonization by staphylococci and gram-negative bacilli (17,90). Although alcohols are among the safest antiseptics available, they can cause dryness and irritation of the skin (1,252). Ethanol is usually less irritating than n-propanol or isopropanol (252).

Irritant contact dermatitis is more commonly reported with iodophors (92). Other antiseptic agents that can cause irritant contact dermatitis (in order of decreasing frequency) include chlorhexidine, PCMX, triclosan, and alcohol-based products. Skin that is damaged by repeated exposure to detergents may be more susceptible to irritation by alcohol-based preparations (253). The irritancy potential of commercially prepared hand-hygiene products, which is often determined by measuring transepidermal water loss, may be available from the manufacturer. Other factors that can contribute to dermatitis associated with frequent handwashing include using hot water for handwashing, low relative humidity (most common in winter months), failure to use supplementary hand lotion or cream, and the quality of paper towels (254,255). Shear forces associated with wearing or removing gloves and allergy to latex proteins may also contribute to dermatitis of the hands of HCWs.

### **Allergic Contact Dermatitis Associated with Hand-Hygiene Products**

Allergic reactions to products applied to the skin (i.e., contact allergies) may present as delayed type reactions (i.e., allergic contact dermatitis) or less commonly as immediate reactions (i.e., contact urticaria). The most common causes of contact allergies are fragrances and preservatives; emulsifiers are less common causes (256–259). Liquid soaps, hand lotions or creams, and “udder ointments” may contain ingredients that cause contact allergies among HCWs (257,258).

Allergic reactions to antiseptic agents, including quaternary ammonium compounds, iodine or iodophors, chlorhexidine, triclosan, PCMX, and alcohols have been reported (118,167,172,256,260–265). Allergic contact dermatitis associated with alcohol-based hand rubs is uncommon. Surveillance at a large hospital in Switzerland, where a commercial alcohol hand rub has been used for >10 years, failed to identify a single case of documented allergy to the product (169). In late 2001, a Freedom of Information Request for data in the FDA’s Adverse Event Reporting System regarding adverse reactions to popular alcohol hand rubs in the United States yielded only one reported case of an erythematous rash reaction attributed to such a product (John M. Boyce, M.D., Hospital of St. Raphael, New Haven, Connecticut, personal communication, 2001). However, with increasing use of such products by HCWs, true allergic reactions to such products likely will be encountered.

Allergic reactions to alcohol-based products may represent true allergy to alcohol, allergy to an impurity or aldehyde metabolite, or allergy to another constituent of the product (167). Allergic contact dermatitis or immediate contact urticarial reactions may be caused by ethanol or isopropanol (167). Allergic reactions can be caused by compounds that may be present as inactive ingredients in alcohol-based hand rubs, including fragrances, benzyl alcohol, stearyl or isostearyl alcohol, phenoxyethanol, myristyl alcohol, propylene glycol, parabens, and benzalkonium chloride (167,256,266–270).

### **Proposed Methods for Reducing Adverse Effects of Agents**

Potential strategies for minimizing hand-hygiene-related irritant contact dermatitis among HCWs include reducing the frequency of exposure to irritating agents (particularly anionic detergents), replacing products with high irritation potential with preparations that cause less damage to the skin, educating personnel regarding the risks of irritant contact dermatitis, and providing caregivers with moisturizing skin-care products or barrier creams (96,98,251,271–273). Reducing the frequency of exposure of HCWs to hand-hygiene products would prove difficult and is not desirable because of the low levels of adherence to hand-hygiene policies in the majority of institutions. Although hospitals have provided personnel with non-antimicrobial soaps in hopes of minimizing dermatitis, frequent use of such products may cause greater skin damage, dryness, and irritation than antiseptic preparations (92,96,98). One strategy for reducing the exposure of personnel to irritating soaps and detergents is to promote the use of alcohol-based hand rubs containing various emollients. Several recent prospective, randomized trials have demonstrated that alcohol-based hand rubs containing emollients were better tolerated by HCWs than washing hands with non-antimicrobial soaps or antimicrobial soaps (96,98,166). Routinely washing hands with soap and water immediately after using an alcohol hand rub may lead to dermatitis. Therefore, personnel should be reminded that it is neither necessary nor recommended to routinely wash hands after each application of an alcohol hand rub.

Hand lotions and creams often contain humectants and various fats and oils that can increase skin hydration and replace altered or depleted skin lipids that contribute to the barrier function of normal skin (251,271). Several controlled trials have demonstrated that regular use (e.g., twice a day) of such products can help prevent and treat irritant contact dermatitis caused by hand-hygiene products (272,273). In one study, frequent and scheduled use of an oil-containing lotion improved skin condition, and thus led to a 50% increase in

handwashing frequency among HCWs (273). Reports from these studies emphasize the need to educate personnel regarding the value of regular, frequent use of hand-care products.

Recently, barrier creams have been marketed for the prevention of hand-hygiene-related irritant contact dermatitis. Such products are absorbed to the superficial layers of the epidermis and are designed to form a protective layer that is not removed by standard handwashing. Two recent randomized, controlled trials that evaluated the skin condition of caregivers demonstrated that barrier creams did not yield better results than did the control lotion or vehicle used (272,273). As a result, whether barrier creams are effective in preventing irritant contact dermatitis among HCWs remains unknown.

In addition to evaluating the efficacy and acceptability of hand-care products, product-selection committees should inquire about the potential deleterious effects that oil-containing products may have on the integrity of rubber gloves and on the efficacy of antiseptic agents used in the facility (8,236).

## Factors To Consider When Selecting Hand-Hygiene Products

When evaluating hand-hygiene products for potential use in health-care facilities, administrators or product-selection committees must consider factors that can affect the overall efficacy of such products, including the relative efficacy of antiseptic agents against various pathogens (Appendix) and acceptance of hand-hygiene products by personnel (274,275). Soap products that are not well-accepted by HCWs can be a deterrent to frequent handwashing (276). Characteristics of a product (either soap or alcohol-based hand rub) that can affect acceptance by personnel include its smell, consistency (i.e., “feel”), and color (92,277,278). For soaps, ease of lathering also may affect user preference.

Because HCWs may wash their hands from a limited number of times per shift to as many as 30 times per shift, the tendency of products to cause skin irritation and dryness is a substantial factor that influences acceptance, and ultimate usage (61,98,274,275,277,279). For example, concern regarding the drying effects of alcohol was a primary cause of poor acceptance of alcohol-based hand-hygiene products in hospitals in the United States (5,143). However, several studies have demonstrated that alcohol-based hand rubs containing emollients are acceptable to HCWs (90,93,98,100,101,106,143,163,164,166). With alcohol-based products, the time required for drying may also affect user acceptance.

Studies indicate that the frequency of handwashing or antiseptic handwashing by personnel is affected by the accessibility of hand-hygiene facilities (280–283). In certain health-care

facilities, only one sink is available in rooms housing several patients, or sinks are located far away from the door of the room, which may discourage handwashing by personnel leaving the room. In intensive-care units, access to sinks may be blocked by bedside equipment (e.g., ventilators or intravenous infusion pumps). In contrast to sinks used for handwashing or antiseptic handwash, dispensers for alcohol-based hand rubs do not require plumbing and can be made available adjacent to each patient’s bed and at many other locations in patient-care areas. Pocket carriage of alcohol-based hand-rub solutions, combined with availability of bedside dispensers, has been associated with substantial improvement in adherence to hand-hygiene protocols (74,284). To avoid any confusion between soap and alcohol hand rubs, alcohol hand-rub dispensers should not be placed adjacent to sinks. HCWs should be informed that washing hands with soap and water after each use of an alcohol hand rub is not necessary and is not recommended, because it may lead to dermatitis. However, because personnel feel a “build-up” of emollients on their hands after repeated use of alcohol hand gels, washing hands with soap and water after 5–10 applications of a gel has been recommended by certain manufacturers.

Automated handwashing machines have not been demonstrated to improve the quality or frequency of handwashing (88,285). Although technologically advanced automated handwashing devices and monitoring systems have been developed recently, only a minimal number of studies have been published that demonstrate that use of such devices results in enduring improvements in hand-hygiene adherence among HCWs. Further evaluation of automated handwashing facilities and monitoring systems is warranted.

Dispenser systems provided by manufacturers or vendors also must be considered when evaluating hand-hygiene products. Dispensers may discourage use by HCWs when they 1) become blocked or partially blocked and do not deliver the product when accessed by personnel, and 2) do not deliver the product appropriately onto the hands. In one hospital where a viscous alcohol-based hand rinse was available, only 65% of functioning dispensers delivered product onto the caregivers’ hands with one press of the dispenser lever, and 9% of dispensers were totally occluded (286). In addition, the volume delivered was often suboptimal, and the product was sometimes squirted onto the wall instead of the caregiver’s hand.

Only limited information is available regarding the cost of hand-hygiene products used in health-care facilities (165,287). These costs were evaluated in patient-care areas at a 450-bed community teaching hospital (287); the hospital spent \$22,000 (\$0.72 per patient-day) on 2% chlorhexidine-containing preparations, plain soap, and an alcohol hand rinse. (287) When

hand-hygiene supplies for clinics and nonpatient care areas were included, the total annual budget for soaps and hand antiseptic agents was \$30,000 (approximately \$1 per patient-day). Annual hand-hygiene product budgets at other institutions vary considerably because of differences in usage patterns and varying product prices. One researcher (287) determined that if non-antimicrobial liquid soap were assigned an arbitrary relative cost of 1.0, the cost per liter would be 1.7 times as much for 2% chlorhexidine gluconate detergent, 1.6–2.0 times higher for alcohol-based hand-rub products, and 4.5 times higher for an alcohol-based foam product. A recent cost comparison of surgical scrubbing with an antimicrobial soap versus brushless scrubbing with an alcohol-based hand rub revealed that costs and time required for preoperative scrubbing were less with the alcohol-based product (165). In a trial conducted in two critical-care units, the cost of using an alcohol hand rub was half as much as using an antimicrobial soap for handwashing (\$0.025 versus \$0.05 per application, respectively) (166).

To put expenditures for hand-hygiene products into perspective, health-care facilities should consider comparing their budget for hand-hygiene products to estimated excess hospital costs resulting from health-care-associated infections. The excess hospital costs associated with only four or five health-care-associated infections of average severity may equal the entire annual budget for hand-hygiene products used in inpatient-care areas. Just one severe surgical site infection, lower respiratory tract infection, or bloodstream infection may cost the hospital more than the entire annual budget for antiseptic agents used for hand hygiene (287). Two studies provided certain quantitative estimates of the benefit of hand-hygiene-promotion programs (72,74). One study demonstrated a cost saving of approximately \$17,000 resulting from reduced use of vancomycin after the observed decrease in MRSA incidence in a 7-month period (72). In another study that examined both direct costs associated with the hand-hygiene promotion program (increased use of hand-rub solution and poster production) and indirect costs associated with health-care-personnel time (74), costs of the program were an estimated \$57,000 or less per year (an average of \$1.42 per patient admitted). Supplementary costs associated with the increased use of alcohol-based hand-rub solution averaged \$6.07 per 100 patient-days. Based on conservative estimates of \$2,100 saved per infection averted and on the assumption that only 25% of the observed reduction in the infection rate was associated with improved hand-hygiene practice, the program was substantially cost-effective. Thus, hospital administrators must consider that by purchasing more effective or more acceptable hand-hygiene products to improve hand-hygiene practices, they

will avoid the occurrence of nosocomial infections; preventing only a limited number of additional health-care-associated infections per year will lead to savings that will exceed any incremental costs of improved hand-hygiene products.

## Hand-Hygiene Practices Among HCWs

In observational studies conducted in hospitals, HCWs washed their hands an average of five times per shift to as many as 30 times per shift (Table 6) (17,61,90,98,274,288); certain nurses washed their hands  $\leq 100$  times per shift (90). Hospitalwide surveillance of hand hygiene reveals that the average number of handwashing opportunities varies markedly between hospital wards. For example, nurses in pediatric wards had an average of eight opportunities for hand hygiene per hour of patient care compared with an average of 20 for nurses in intensive-care units (11). The duration of handwashing or hygienic handwash episodes by HCWs has averaged 6.6–24.0 seconds in observational studies (Table 7) (17,52,59,84–87,89,249,279). In addition to washing their

**TABLE 6. Handwashing frequency among health-care workers**

| Ref. no. | Year | Avg. no./time period | Range  | Avg. no./hr |
|----------|------|----------------------|--------|-------------|
| (61)     | 1988 | 5/8 hour             | N.S.   |             |
| (89)     | 1984 | 5–10/shift           | N.S.   |             |
| (96)     | 2000 | 10/shift             | N.S.   |             |
| (273)    | 2000 | 12–18/day            | 2–60   |             |
| (98)     | 2000 | 13–15/8 hours        | 5–27   | 1.6–1.8/hr  |
| (90)     | 1977 | 20–42/8 hours        | 10–100 |             |
| (391)    | 2000 | 21/12 hours          | N.S.   |             |
| (272)    | 2000 | 22/day               | 0–70   |             |
| (88)     | 1991 |                      |        | 1.7–2.1/hr  |
| (17)     | 1998 |                      |        | 2.1/hr      |
| (279)    | 1978 |                      |        | 3/hr        |
| (303)    | 1994 |                      |        | 3.3/hr      |

Note: N.S. = Not Stated.

**TABLE 7. Average duration of handwashing by health-care workers**

| Ref. no. | Year | Mean/median time  |
|----------|------|-------------------|
| (392)    | 1997 | 4.7–5.3 seconds   |
| (303)    | 1994 | 6.6 seconds       |
| (52)     | 1974 | 8–9.3 seconds     |
| (85)     | 1984 | 8.6 seconds       |
| (86)     | 1994 | <9 seconds        |
| (87)     | 1994 | 9.5 seconds       |
| (88)     | 1991 | <10 seconds       |
| (294)    | 1990 | 10 seconds        |
| (89)     | 1984 | 11.6 seconds      |
| (300)    | 1992 | 12.5 seconds      |
| (59)     | 1988 | 15.6–24.4 seconds |
| (17)     | 1998 | 20.6 seconds      |
| (279)    | 1978 | 21 seconds        |
| (293)    | 1989 | 24 seconds        |

hands for limited time periods, personnel often fail to cover all surfaces of their hands and fingers (288).

### Adherence of HCWs to Recommended Hand-Hygiene Practices

Observational Studies of Hand-Hygiene Adherence. Adherence of HCWs to recommended hand-hygiene procedures has been poor, with mean baseline rates of 5%–81% (overall average: 40%) (Table 8) (71,74,86,87,276,280,281,283,285,289–313). The methods used for defining adherence (or non-adherence) and those used for conducting observations vary considerably among studies, and reports do not provide

detailed information concerning the methods and criteria used. The majority of studies were conducted with hand-hygiene adherence as the major outcome measure, whereas a limited number measured adherence as part of a broader investigation. Several investigators reported improved adherence after implementing various interventions, but the majority of studies had short follow-up periods and did not confirm whether behavioral improvements were long-lasting. Other studies established that sustained improvements in handwashing behavior occurred during a long-term program to improve adherence to hand-hygiene policies (74,75).

**TABLE 8. Hand-hygiene adherence by health-care workers (1981–2000)**

| Ref. no. | Year | Setting          | Before/after | Adherence baseline | Adherence after intervention | Intervention   |
|----------|------|------------------|--------------|--------------------|------------------------------|--|
| (280)    | 1981 | ICU              | A            | 16%                | 30%                          | More convenient sink locations   |
| (289)    | 1981 | ICU              | A            | 41%                | —                            |  |
|          |      | ICU              | A            | 28%                | —                            |  |
| (290)    | 1983 | All wards        | A            | 45%                | —                            |  |
| (281)    | 1986 | SICU             | A            | 51%                | —                            |  |
|          |      | MICU             | A            | 76%                | —                            |  |
| (276)    | 1986 | ICU              | A            | 63%                | 92%                          | Performance feedback   |
| (291)    | 1987 | PICU             | A            | 31%                | 30%                          | Wearing overgown   |
| (292)    | 1989 | MICU             | B/A          | 14%/28%*           | 73%/81%                      | Feedback, policy reviews, memo, and posters                                  |
|          |      | MICU             | B/A          | 26%/23%            | 38%/60%                      |  |
| (293)    | 1989 | NICU             | A/B          | 75%/50%            | —                            |  |
| (294)    | 1990 | ICU              | A            | 32%                | 45%                          | Alcohol rub introduced   |
| (295)    | 1990 | ICU              | A            | 81%                | 92%                          | Inservices first, then group feedback  |
| (296)    | 1990 | ICU              | B/A          | 22%                | 30%                          |  |
| (297)    | 1991 | SICU             | A            | 51%                | —                            |  |
| (298)    | 1991 | Pedi OPDs        | B            | 49%                | 49%                          | Signs, feedback, and verbal reminders to physicians                          |
| (299)    | 1991 | Nursery and NICU | B/A†         | 28%                | 63%                          | Feedback, dissemination of literature, and results of environmental cultures |
| (300)    | 1992 | NICU/others      | A            | 29%                | —                            |  |
| (71)     | 1992 | ICU              | N.S.         | 40%                | —                            |  |
| (301)    | 1993 | ICUs             | A            | 40%                | —                            |  |
| (87)     | 1994 | Emergency Room   | A            | 32%                | —                            |  |
| (86)     | 1994 | All wards        | A            | 32%                | —                            |  |
| (285)    | 1994 | SICU             | A            | 22%                | 38%                          | Automated handwashing machines available                                     |
| (302)    | 1994 | NICU             | A            | 62%                | 60%                          | No gowning required  |
| (303)    | 1994 | ICU Wards        | AA           | 30%/29%            | —                            |  |
| (304)    | 1995 | ICU Oncol Ward   | A            | 56%                | —                            |  |
| (305)    | 1995 | ICU              | N.S.         | 5%                 | 63%                          | Lectures, feedback, and demonstrations                                       |
| (306)    | 1996 | PICU             | B/A          | 12%/11%            | 68%/65%                      | Overt observation, followed by feedback                                      |
| (307)    | 1996 | MICU             | A            | 41%                | 58%                          | Routine wearing of gowns and gloves  |
| (308)    | 1996 | Emergency Dept   | A            | 54%                | 64%                          | Signs/distributed review paper   |
| (309)    | 1998 | All wards        | A            | 30%                | —                            |  |
| (310)    | 1998 | Pediatric wards  | B/A          | 52%/49%            | 74%/69%                      | Feedback, movies, posters, and brochures                                     |
| (311)    | 1999 | MICU             | B/A          | 12%/55%            | —                            |  |
| (74)     | 2000 | All wards        | B/A          | 48%                | 67%                          | Posters, feedback, administrative support, and alcohol rub                   |
| (312)    | 2000 | MICU             | A            | 42%                | 61%                          | Alcohol hand rub made available  |
| (283)    | 2000 | MICU             | B/A          | 10%/22%            | 23%/48%                      | Education, feedback, and alcohol gel made available                          |
|          |      | CTICU            | B/A          | 4%/13%             | 7%/14%                       |  |
| (313)    | 2000 | Medical wards    | A            | 60%                | 52%                          | Education, reminders, and alcohol gel made available                         |

**Note:** ICU = intensive care unit, SICU = surgical ICU, MICU = medical ICU, PICU = pediatric ICU, NICU = neonatal ICU, Emerg = emergency, Oncol = oncology, CTICU = cardiothoracic ICU, and N.S. = not stated.

\* Percentage compliance before/after patient contact.

† After contact with inanimate objects.



**Factors Affecting Adherence.** Factors that may influence hand hygiene include those identified in epidemiologic studies and factors reported by HCWs as being reasons for lack of adherence to hand-hygiene recommendations. Risk factors for poor adherence to hand hygiene have been determined objectively in several observational studies or interventions to improve adherence (11,12,274,292,295,314–317). Among these, being a physician or a nursing assistant, rather than a nurse, was consistently associated with reduced adherence (Box 1).

In the largest hospitalwide survey of hand-hygiene practices among HCWs (11), predictors of poor adherence to recommended hand-hygiene measures were identified. Predictor variables included professional category, hospital ward, time of day/week, and type and intensity of patient care, defined as the number of opportunities for hand hygiene per hour of patient care. In 2,834 observed opportunities for hand hygiene, average adherence was 48%. In multivariate analysis, nonadherence was lowest among nurses and during weekends

### BOX 1. Factors influencing adherence to hand-hygiene practices\*

#### Observed risk factors for poor adherence to recommended hand-hygiene practices

- Physician status (rather than a nurse)
- Nursing assistant status (rather than a nurse)
- Male sex
- Working in an intensive-care unit
- Working during the week (versus the weekend)
- Wearing gowns/gloves
- Automated sink
- Activities with high risk of cross-transmission
- High number of opportunities for hand hygiene per hour of patient care

#### Self-reported factors for poor adherence with hand hygiene

- Handwashing agents cause irritation and dryness
- Sinks are inconveniently located/shortage of sinks
- Lack of soap and paper towels
- Often too busy/insufficient time
- Understaffing/overcrowding
- Patient needs take priority
- Hand hygiene interferes with health-care worker relationships with patients
- Low risk of acquiring infection from patients
- Wearing of gloves/beliefs that glove use obviates the need for hand hygiene
- Lack of knowledge of guidelines/protocols
- Not thinking about it/forgetfulness
- No role model from colleagues or superiors
- Skepticism regarding the value of hand hygiene
- Disagreement with the recommendations
- Lack of scientific information of definitive impact of improved hand hygiene on health-care-associated infection rates

#### Additional perceived barriers to appropriate hand hygiene

- Lack of active participation in hand-hygiene promotion at individual or institutional level
- Lack of role model for hand hygiene
- Lack of institutional priority for hand hygiene
- Lack of administrative sanction of noncompliers/rewarding compliers
- Lack of institutional safety climate

\* Source: Adapted from Pittet D. Improving compliance with hand hygiene in hospitals. *Infect Control Hosp Epidemiol* 2000;21:381–6.

(Odds Ratio [OR]: 0.6; 95% confidence interval [CI] = 0.4–0.8). Nonadherence was higher in intensive-care units compared with internal medicine wards (OR: 2.0; 95% CI = 1.3–3.1), during procedures that carried a high risk of bacterial contamination (OR: 1.8; 95% CI = 1.4–2.4), and when intensity of patient care was high (21–40 handwashing opportunities — OR: 1.3; 95% CI = 1.0–1.7; 41–60 opportunities — OR: 2.1; 95% CI = 1.5–2.9; >60 opportunities — OR: 2.1; 95% CI = 1.3–3.5). The higher the demand for hand hygiene, the lower the adherence; on average, adherence decreased by 5% ( $\pm$  2%) for each increase of 10 opportunities per hour when the intensity of patient care exceeded 10 opportunities per hour. Similarly, the lowest adherence rate (36%) was found in intensive-care units, where indications for hand hygiene were typically more frequent (on average, 20 opportunities per patient-hour). The highest adherence rate (59%) was observed in pediatrics wards, where the average intensity of patient care was lower than in other hospital areas (an average of eight opportunities per patient-hour). The results of this study indicate that full adherence to previous guidelines may be unrealistic, and that facilitated access to hand hygiene could help improve adherence (11,12,318).

Perceived barriers to adherence with hand-hygiene practice recommendations include skin irritation caused by hand-hygiene agents, inaccessible hand-hygiene supplies, interference with HCW-patient relationships, priority of care (i.e., the patients' needs are given priority over hand hygiene), wearing of gloves, forgetfulness, lack of knowledge of the guidelines, insufficient time for hand hygiene, high workload and understaffing, and the lack of scientific information indicating a definitive impact of improved hand hygiene on health-care-associated infection rates (11,274,292,295,315–317). Certain perceived barriers to adherence with hand-hygiene guidelines have been assessed or quantified in observational studies (12,274,292,295,314–317) (Box 1).

Skin irritation by hand-hygiene agents constitutes a substantial barrier to appropriate adherence (319). Because soaps and detergents can damage skin when applied on a regular basis, HCWs must be better informed regarding the possible adverse effects associated with hand-hygiene agents. Lack of knowledge and education regarding this subject is a barrier to motivation. In several studies, alcohol-based hand rubs containing emollients (either isopropanol, ethanol, or n-propanol in 60%–90% vol/vol) were less irritating to the skin than the soaps or detergents tested. In addition, the alcohol-based products containing emollients that were tested were at least as tolerable and efficacious as the detergents tested. Also, studies demonstrate that several hand lotions have reduced skin scaling and cracking, which may reduce microbial shedding from the hands (67,272,273).

Easy access to hand-hygiene supplies, whether sink, soap, medicated detergent, or alcohol-based hand-rub solution, is essential for optimal adherence to hand-hygiene recommendations. The time required for nurses to leave a patient's bedside, go to a sink, and wash and dry their hands before attending the next patient is a deterrent to frequent handwashing or hand antisepsis (11,318). Engineering controls could facilitate adherence, but careful monitoring of hand-hygiene behavior should be conducted to exclude the possible negative effect of newly introduced handwashing devices (88).

The impact of wearing gloves on adherence to hand-hygiene policies has not been definitively established, because published studies have yielded contradictory results (87,290,301,320). Hand hygiene is required regardless of whether gloves are used or changed. Failure to remove gloves after patient contact or between "dirty" and "clean" body-site care on the same patient must be regarded as nonadherence to hand-hygiene recommendations (11). In a study in which experimental conditions approximated those occurring in clinical practice (321), washing and reusing gloves between patient contacts resulted in observed bacterial counts of 0–4.7 log on the hands after glove removal. Therefore, this practice should be discouraged; handwashing or disinfection should be performed after glove removal.

Lack of 1) knowledge of guidelines for hand hygiene, 2) recognition of hand-hygiene opportunities during patient care, and 3) awareness of the risk of cross-transmission of pathogens are barriers to good hand-hygiene practices. Furthermore, certain HCWs believe they have washed their hands when necessary, even when observations indicate they have not (89,92,295,296,322).

Perceived barriers to hand-hygiene behavior are linked not only to the institution, but also to HCWs' colleagues. Therefore, both institutional and small-group dynamics need to be considered when implementing a system change to secure an improvement in HCWs' hand-hygiene practice.

### Possible Targets for Hand-Hygiene Promotion

Targets for the promotion of hand hygiene are derived from studies assessing risk factors for nonadherence, reported reasons for the lack of adherence to recommendations, and additional factors perceived as being important to facilitate appropriate HCW behavior. Although certain factors cannot be modified (Box 1), others can be changed.

One factor that must be addressed is the time required for HCWs to clean their hands. The time required for traditional handwashing may render full adherence to previous guidelines unrealistic (11,12,318) and more rapid access to hand-hygiene materials could help improve adherence. One study conducted in an intensive-care unit demonstrated that it took

nurses an average of 62 seconds to leave a patient's bedside, walk to a sink, wash their hands, and return to patient care (318). In contrast, an estimated one fourth as much time is required when using alcohol-based hand rub placed at each patient's bedside. Providing easy access to hand-hygiene materials is mandatory for appropriate hand-hygiene behavior and is achievable in the majority of health-care facilities (323). In particular, in high-demand situations (e.g., the majority of critical-care units), under hectic working conditions, and at times of overcrowding or understaffing, HCWs may be more likely to use an alcohol-based hand rub than to wash their hands (323). Further, using alcohol-based hand rubs may be a better option than traditional handwashing with plain soap and water or antiseptic handwash, because they not only require less time (166,318) but act faster (1) and irritate hands less often (1,67,96,98,166). They also were used in the only program that reported a sustained improvement in hand-hygiene adherence associated with decreased infection rates (74). However, making an alcohol-based hand rub available to personnel without providing ongoing educational and motivational activities may not result in long-lasting improvement in hand-hygiene practices (313). Because increased use of hand-hygiene agents might be associated with skin dryness, the availability of free skin-care lotion is recommended.

Education is a cornerstone for improvement with hand-hygiene practices. Topics that must be addressed by educational programs include the lack of 1) scientific information for the definitive impact of improved hand hygiene on health-care-associated infection and resistant organism transmission rates; 2) awareness of guidelines for hand hygiene and insufficient knowledge concerning indications for hand hygiene during daily patient care; 3) knowledge concerning the low average adherence rate to hand hygiene by the majority of HCWs; and 4) knowledge concerning the appropriateness, efficacy, and understanding of the use of hand-hygiene and skin-care-protection agents.

HCWs necessarily evolve within a group that functions within an institution. Possible targets for improvement in hand-hygiene behavior not only include factors linked to individual HCWs, but also those related to the group(s) and the institution as a whole (317,323). Examples of possible targets for hand-hygiene promotion at the group level include education and performance feedback on hand-hygiene adherence; efforts to prevent high workload, downsizing, and understaffing; and encouragement and provision of role models from key members in the work unit. At the institutional level, targets for improvement include 1) written guidelines, hand-hygiene agents, skin-care promotions and agents, or hand-hygiene facilities; 2) culture or tradition of adherence; and 3)

administrative leadership, sanction, support, and rewards. Several studies, conducted in various types of institutions, reported modest and even low levels of adherence to recommended hand-hygiene practices, indicating that such adherence varied by hospital ward and by type of HCW. These results indicate educational sessions may need to be designed specifically for certain types of personnel (11,289,290,294,317,323).

## Lessons Learned from Behavioral Theories

In 1998, the prevailing behavioral theories and their applications with regard to the health professions were reviewed by researchers in an attempt to better understand how to target more successful interventions (317). The researchers proposed a hypothetical framework to enhance hand-hygiene practices and stressed the importance of considering the complexity of individual and institutional factors when designing behavioral interventions.

Although behavioral theories and secondary interventions have primarily targeted individual workers, this practice might be insufficient to produce sustained change (317,324,325). Interventions aimed at improving hand-hygiene practices must account for different levels of behavior interaction (12,317,326). Thus, the interdependence of individual factors, environmental constraints, and the institutional climate must be taken into account in the strategic planning and development of hand-hygiene campaigns. Interventions to promote hand hygiene in hospitals should consider variables at all these levels. Various factors involved in hand-hygiene behavior include intention, attitude towards the behavior, perceived social norm, perceived behavioral control, perceived risk for infection, hand-hygiene practices, perceived role model, perceived knowledge, and motivation (317). The factors necessary for change include 1) dissatisfaction with the current situation, 2) perception of alternatives, and 3) recognition, both at the individual and institutional level, of the ability and potential to change. Although the latter implies education and motivation, the former two necessitate a system change.

Among the reported reasons for poor adherence with hand-hygiene recommendations (Box 1), certain ones are clearly associated with the institution or system (e.g., lack of institutional priority for hand hygiene, administrative sanctions, and a safety climate). Although all of these reasons would require a system change in the majority of institutions, the third requires management commitment, visible safety programs, an acceptable level of work stress, a tolerant and supportive attitude toward reported problems, and belief in the efficacy

of preventive strategies (12,317,325,327). Most importantly, an improvement in infection-control practices requires 1) questioning basic beliefs, 2) continuous assessment of the group (or individual) stage of behavioral change, 3) intervention(s) with an appropriate process of change, and 4) supporting individual and group creativity (317). Because of the complexity of the process of change, single interventions often fail. Thus, a multimodal, multidisciplinary strategy is likely necessary (74,75,317,323,326).

## Methods Used To Promote Improved Hand Hygiene

Hand-hygiene promotion has been challenging for >150 years. In-service education, information leaflets, workshops and lectures, automated dispensers, and performance feedback on hand-hygiene adherence rates have been associated with transient improvement (291,294–296,306,314).

Several strategies for promotion of hand hygiene in hospitals have been published (Table 9). These strategies require education, motivation, or system change. Certain strategies are based on epidemiologic evidence, others on the authors' and other investigators' experience and review of current knowledge. Some strategies may be unnecessary in certain circumstances, but may be helpful in others. In particular, changing the hand-hygiene agent could be beneficial in institutions or hospital wards with a high workload and a high demand for hand hygiene when alcohol-based hand rubs are not available (11,73,78,328). However, a change in the recommended hand-hygiene agent could be deleterious if introduced during winter, at a time of higher hand-skin irritability, and if not accompanied by the provision of skin-care products (e.g., pro-

TECTIVE creams and lotions). Additional specific elements should be considered for inclusion in educational and motivational programs (Box 2).

Several strategies that could potentially be associated with successful promotion of hand hygiene require a system change (Box 1). Hand-hygiene adherence and promotion involve factors at both the individual and system level. Enhancing individual and institutional attitudes regarding the feasibility of making changes (self-efficacy), obtaining active participation of personnel at both levels, and promoting an institutional safety climate represent challenges that exceed the current perception of the role of infection-control professionals.

Whether increased education, individual reinforcement technique, appropriate rewarding, administrative sanction, enhanced self-participation, active involvement of a larger number of organizational leaders, enhanced perception of health threat, self-efficacy, and perceived social pressure (12,317,329,330), or combinations of these factors can improve HCWs' adherence with hand hygiene needs further investigation. Ultimately, adherence to recommended hand-hygiene practices should become part of a culture of patient safety where a set of interdependent quality elements interact to achieve a shared objective (331).

On the basis of both these hypothetical considerations and successful, actual experiences in certain institutions, strategies to improve adherence to hand-hygiene practices should be both multimodal and multidisciplinary. However, strategies must be further researched before they are implemented.

**TABLE 9. Strategies for successful promotion of hand hygiene in hospitals**

| Strategy  | Tool for change* | Selected references†    |
|---|------------------|-------------------------|
| Education   | E (M, S)         | (74,295,306,326,393)    |
| Routine observation and feedback                                  | S (E, M)         | (74,294,306,326,393)    |
| Engineering control   |                  |                         |
| Make hand hygiene possible, easy, and convenient                  | S                | (74,281,326,393)        |
| Make alcohol-based hand rub available                             | S                | (74)                    |
| (at least in high-demand situations)                              | S                | (74,283,312)            |
| Patient education   | S (M)            | (283,394)               |
| Reminders in the workplace  | S                | (74,395)                |
| Administrative sanction/rewarding                                 | S                | (12,317)                |
| Change in hand-hygiene agent                                      | S (E)            | (11,67,71,283,312)      |
| Promote/facilitate skin care for health-care-workers' hands       | S (E)            | (67,74,274,275)         |
| Obtain active participation at individual and institutional level | E, M, S          | (74,75,317)             |
| Improve institutional safety climate                              | S (M)            | (74,75,317)             |
| Enhance individual and institutional self-efficacy                | S (E, M)         | (74,75,317)             |
| Avoid overcrowding, understaffing, and excessive workload         | S                | (11,74,78,297,396)      |
| Combine several of above strategies                               | E, M, S          | (74,75,295,306,317,326) |

\* The dynamic of behavioral change is complex and involves a combination of education (E), motivation (M), and system change (S).

† Only selected references have been listed; readers should refer to more extensive reviews for exhaustive reference lists (1,8,317,323,397).



**BOX 2. Elements of health-care worker educational and motivational programs****Rationale for hand hygiene**

- Potential risks of transmission of microorganisms to patients
- Potential risks of health-care worker colonization or infection caused by organisms acquired from the patient
- Morbidity, mortality, and costs associated with health-care-associated infections

**Indications for hand hygiene**

- Contact with a patient's intact skin (e.g., taking a pulse or blood pressure, performing physical examinations, lifting the patient in bed) (25,26,45,48,51,53)
- Contact with environmental surfaces in the immediate vicinity of patients (46,51,53,54)
- After glove removal (50,58,71)

**Techniques for hand hygiene**

- Amount of hand-hygiene solution
- Duration of hand-hygiene procedure
- Selection of hand-hygiene agents
  - Alcohol-based hand rubs are the most efficacious agents for reducing the number of bacteria on the hands of personnel. Antiseptic soaps and detergents are the next most effective, and non-antimicrobial soaps are the least effective (1,398).
  - Soap and water are recommended for visibly soiled hands.
  - Alcohol-based hand rubs are recommended for routine decontamination of hands for all clinical indications (except when hands are visibly soiled) and as one of the options for surgical hand hygiene.

**Methods to maintain hand skin health**

- Lotions and creams can prevent or minimize skin dryness and irritation caused by irritant contact dermatitis
- Acceptable lotions or creams to use
- Recommended schedule for applying lotions or creams

**Expectations of patient care managers/administrators**

- Written statements regarding the value of, and support for, adherence to recommended hand-hygiene practices
- Role models demonstrating adherence to recommended hand hygiene practices (399)

**Indications for, and limitations of, glove use**

- Hand contamination may occur as a result of small, undetected holes in examination gloves (321,361)
- Contamination may occur during glove removal (50)
- Wearing gloves does not replace the need for hand hygiene (58)
- Failure to remove gloves after caring for a patient may lead to transmission of microorganisms from one patient to another (373).

**Efficacy of Promotion and Impact of Improved Hand Hygiene**

The lack of scientific information of the definitive impact of improved hand hygiene on health-care-associated infection rates is a possible barrier to appropriate adherence with hand-hygiene recommendations (Box 1). However, evidence supports the belief that improved hand hygiene can reduce health-care-associated infection rates. Failure to perform appropriate hand hygiene is considered the leading cause of

health-care-associated infections and spread of multiresistant organisms and has been recognized as a substantial contributor to outbreaks.

Of nine hospital-based studies of the impact of hand hygiene on the risk of health-care-associated infections (Table 10) (48,69–75,296), the majority demonstrated a temporal relationship between improved hand-hygiene practices and reduced infection rates.

In one of these studies, endemic MRSA in a neonatal intensive-care unit was eliminated 7 months after introduction of a new

**TABLE 10. Association between improved adherence with hand-hygiene practice and health-care–associated infection rates**

| Year | Ref. no. | Hospital setting | Results   | Duration of follow-up |
|------|----------|------------------|---|-----------------------|
| 1977 | (48)     | Adult ICU        | Reduction in health-care–associated infections caused by endemic <i>Klebsiella</i> spp.   | 2 years               |
| 1982 | (69)     | Adult ICU        | Reduction in health-care-associated infection rates   | N.S.                  |
| 1984 | (70)     | Adult ICU        | Reduction in health-care–associated infection rates   | N.S.                  |
| 1990 | (296)    | Adult ICU        | No effect (average hand hygiene adherence improvement did not reach statistical significance)   | 11 months             |
| 1992 | (71)     | Adult ICU        | Substantial difference between rates of health-care–associated infection between two different hand-hygiene agents  | 8 months              |
| 1994 | (72)     | NICU             | Elimination of MRSA, when combined with multiple other infection-control measures. Reduction of vancomycin use  | 9 months              |
| 1995 | (73)     | Newborn nursery  | Elimination of MRSA, when combined with multiple other infection-control measures   | 3.5 years             |
| 2000 | (75)     | MICU/NICU        | 85% relative reduction of VRE rate in the intervention hospital; 44% relative reduction in control hospital; no change in MRSA  | 8 months              |
| 2000 | (74)     | Hospitalwide     | Substantial reduction in the annual overall prevalence of health-care–associated infections and MRSA cross-transmission rates. Active surveillance cultures and contact precautions were implemented during same period | 5 years               |

**Note:** ICU = intensive care unit, NICU = neonatal ICU, MRSA = methicillin-resistant *Staphylococcus aureus*, MICU = medical ICU, and N.S. = not stated.

hand antiseptic (1% triclosan); all other infection-control measures remained in place, including the practice of conducting weekly active surveillance by obtaining cultures (72). Another study reported an MRSA outbreak involving 22 infants in a neonatal unit (73). Despite intensive efforts, the outbreak could not be controlled until a new antiseptic was added (i.e., 0.3% triclosan); all previously used control measures remained in place, including gloves and gowns, cohorting, and obtaining cultures for active surveillance.

The effectiveness of a longstanding, hospitalwide program to promote hand hygiene at the University of Geneva hospitals was recently reported (74). Overall adherence to hand-hygiene guidelines during routine patient care was monitored during hospitalwide observational surveys. These surveys were conducted biannually during December 1994–December 1997, before and during implementation of a hand-hygiene campaign that specifically emphasized the practice of bedside, alcohol-based hand disinfection. Individual-sized bottles of hand-rub solution were distributed to all wards, and custom-made holders were mounted on all beds to facilitate access to hand disinfection. HCWs were also encouraged to carry bottles in their pockets, and in 1996, a newly designed flat (instead of round) bottle was made available to further facilitate pocket carriage. The promotional strategy was multimodal and involved a multidisciplinary team of HCWs, the use of wall posters, the promotion of antiseptic hand rubs located at bed-sides throughout the institution, and regular performance feedback to all HCWs (see <http://www.hopisafe.ch> for further

details on methodology). Health-care–associated infection rates, attack rates of MRSA cross-transmission, and consumption of hand-rub disinfectant were measured. Adherence to recommended hand-hygiene practices improved progressively from 48% in 1994 to 66% in 1997 ( $p < 0.001$ ). Whereas recourse to handwashing with soap and water remained stable, frequency of hand disinfection markedly increased during the study period ( $p < 0.001$ ), and the consumption of alcohol-based hand-rub solution increased from 3.5 to 15.4 liters per 1,000 patient-days during 1993–1998 ( $p < 0.001$ ). The increased frequency of hand disinfection was unchanged after adjustment for known risk factors of poor adherence. During the same period, both overall health-care–associated infection and MRSA transmission rates decreased (both  $p < 0.05$ ). The observed reduction in MRSA transmission may have been affected by both improved hand-hygiene adherence and the simultaneous implementation of active surveillance cultures for detecting and isolating patients colonized with MRSA (332). The experience from the University of Geneva hospitals constitutes the first report of a hand-hygiene campaign with a sustained improvement over several years. An additional multimodal program also yielded sustained improvements in hand-hygiene practices over an extended period (75); the majority of studies have been limited to a 6- to 9-month observation period.

Although these studies were not designed to assess the independent contribution of hand hygiene on the prevention of health-care–associated infections, the results indicate that

improved hand-hygiene practices reduce the risk of transmission of pathogenic microorganisms. The beneficial effects of hand-hygiene promotion on the risk of cross-transmission also have been reported in surveys conducted in schools and day care centers (333–338), as well as in a community setting (339–341).

## Other Policies Related to Hand Hygiene

### Fingernails and Artificial Nails

Studies have documented that subungual areas of the hand harbor high concentrations of bacteria, most frequently coagulase-negative staphylococci, gram-negative rods (including *Pseudomonas* spp.), Corynebacteria, and yeasts (14,342,343). Freshly applied nail polish does not increase the number of bacteria recovered from periungual skin, but chipped nail polish may support the growth of larger numbers of organisms on fingernails (344,345). Even after careful handwashing or the use of surgical scrubs, personnel often harbor substantial numbers of potential pathogens in the subungual spaces (346–348).

Whether artificial nails contribute to transmission of health-care-associated infections is unknown. However, HCWs who wear artificial nails are more likely to harbor gram-negative pathogens on their fingertips than are those who have natural nails, both before and after handwashing (347–349). Whether the length of natural or artificial nails is a substantial risk factor is unknown, because the majority of bacterial growth occurs along the proximal 1 mm of the nail adjacent to subungual skin (345,347,348). Recently, an outbreak of *P. aeruginosa* in a neonatal intensive care unit was attributed to two nurses (one with long natural nails and one with long artificial nails) who carried the implicated strains of *Pseudomonas* spp. on their hands (350). Patients were substantially more likely than controls to have been cared for by the two nurses during the exposure period, indicating that colonization of long or artificial nails with *Pseudomonas* spp. may have contributed to causing the outbreak. Personnel wearing artificial nails also have been epidemiologically implicated in several other outbreaks of infection caused by gram-negative bacilli and yeast (351–353). Although these studies provide evidence that wearing artificial nails poses an infection hazard, additional studies are warranted.

### Gloving Policies

CDC has recommended that HCWs wear gloves to 1) reduce the risk of personnel acquiring infections from patients, 2) prevent health-care worker flora from being transmitted to patients, and 3) reduce transient contamination of the hands

of personnel by flora that can be transmitted from one patient to another (354). Before the emergence of the acquired immunodeficiency syndrome (AIDS) epidemic, gloves were worn primarily by personnel caring for patients colonized or infected with certain pathogens or by personnel exposed to patients with a high risk of hepatitis B. Since 1987, a dramatic increase in glove use has occurred in an effort to prevent transmission of HIV and other bloodborne pathogens from patients to HCWs (355). The Occupational Safety and Health Administration (OSHA) mandates that gloves be worn during all patient-care activities that may involve exposure to blood or body fluids that may be contaminated with blood (356).

The effectiveness of gloves in preventing contamination of HCWs' hands has been confirmed in several clinical studies (45,51,58). One study found that HCWs who wore gloves during patient contact contaminated their hands with an average of only 3 CFUs per minute of patient care, compared with 16 CFUs per minute for those not wearing gloves (51). Two other studies, involving personnel caring for patients with *C. difficile* or VRE, revealed that wearing gloves prevented hand contamination among the majority of personnel having direct contact with patients (45,58). Wearing gloves also prevented personnel from acquiring VRE on their hands when touching contaminated environmental surfaces (58). Preventing heavy contamination of the hands is considered important, because handwashing or hand antisepsis may not remove all potential pathogens when hands are heavily contaminated (25,111).

Several studies provide evidence that wearing gloves can help reduce transmission of pathogens in health-care settings. In a prospective controlled trial that required personnel to routinely wear vinyl gloves when handling any body substances, the incidence of *C. difficile* diarrhea among patients decreased from 7.7 cases/1,000 patient discharges before the intervention to 1.5 cases/1,000 discharges during the intervention (226). The prevalence of asymptomatic *C. difficile* carriage also decreased substantially on "glove" wards, but not on control wards. In intensive-care units where VRE or MRSA have been epidemic, requiring all HCWs to wear gloves to care for all patients in the unit (i.e., universal glove use) likely has helped control outbreaks (357,358).

The influence of glove use on the hand-hygiene habits of personnel is not clear. Several studies found that personnel who wore gloves were less likely to wash their hands upon leaving a patient's room (290,320). In contrast, two other studies found that personnel who wore gloves were substantially more likely to wash their hands after patient care (87,301).

The following caveats regarding use of gloves by HCWs must be considered. Personnel should be informed that gloves

do not provide complete protection against hand contamination. Bacterial flora colonizing patients may be recovered from the hands of  $\leq 30\%$  of HCWs who wear gloves during patient contact (50,58). Further, wearing gloves does not provide complete protection against acquisition of infections caused by hepatitis B virus and herpes simplex virus (359,360). In such instances, pathogens presumably gain access to the caregiver's hands via small defects in gloves or by contamination of the hands during glove removal (50,321,359,361).

Gloves used by HCWs are usually made of natural rubber latex and synthetic nonlatex materials (e.g., vinyl, nitrile, and neoprene [polymers and copolymers of chloroprene]). Because of the increasing prevalence of latex sensitivity among HCWs and patients, FDA has approved several powdered and powder-free latex gloves with reduced protein contents, as well as synthetic gloves that can be made available by health-care institutions for use by latex-sensitive employees. In published studies, the barrier integrity of gloves varies on the basis of type and quality of glove material, intensity of use, length of time used, manufacturer, whether gloves were tested before or after use, and method used to detect glove leaks (359,361–366). In published studies, vinyl gloves have had defects more frequently than latex gloves, the difference in defect frequency being greatest after use (359,361,364,367). However, intact vinyl gloves provide protection comparable to that of latex gloves (359). Limited studies indicate that nitrile gloves have leakage rates that approximate those of latex gloves (368–371). Having more than one type of glove available is desirable, because it allows personnel to select the type that best suits their patient-care activities. Although recent studies indicate that improvements have been made in the quality of gloves (366), hands should be decontaminated or washed after removing gloves (8,50,58,321,361). Gloves should not be washed or reused (321,361). Use of petroleum-based hand lotions or creams may adversely affect the integrity of latex gloves (372). After use of powdered gloves, certain alcohol hand rubs may interact with residual powder on the hands of personnel, resulting in a gritty feeling on the hands. In facilities where powdered gloves are commonly used, various alcohol-based hand rubs should be tested after removal of powdered gloves to avoid selecting a product that causes this undesirable reaction. Personnel should be reminded that failure to remove gloves between patients may contribute to transmission of organisms (358,373).

## Jewelry

Several studies have demonstrated that skin underneath rings is more heavily colonized than comparable areas of skin on fingers without rings (374–376). One study found that 40% of nurses harbored gram-negative bacilli (e.g., *E. cloacae*, *Klebsiella*, and *Acinetobacter*) on skin under rings and that certain nurses carried the same organism under their rings for several months (375). In a more recent study involving >60 intensive care unit nurses, multivariable analysis revealed that rings were the only substantial risk factor for carriage of gram-negative bacilli and *S. aureus* and that the concentration of organisms recovered correlated with the number of rings worn (377). Whether the wearing of rings results in greater transmission of pathogens is unknown. Two studies determined that mean bacterial colony counts on hands after handwashing were similar among persons wearing rings and those not wearing rings (376,378). Further studies are needed to establish if wearing rings results in greater transmission of pathogens in health-care settings.

## Hand-Hygiene Research Agenda

Although the number of published studies concerning hand hygiene has increased considerably in recent years, many questions regarding hand-hygiene products and strategies for improving adherence of personnel to recommended policies remain unanswered. Several concerns must still be addressed by researchers in industry and by clinical investigators (Box 3).

## Web-Based Hand-Hygiene Resources

Additional information regarding improving hand hygiene is available at <http://www.hopisafe.ch>

University of Geneva Hospitals, Geneva, Switzerland

<http://www.cdc.gov/ncidod/hip>

CDC, Atlanta, Georgia

<http://www.jr2.ox.ac.uk/bandolier/band88/b88-8.html>

Bandolier journal, United Kingdom

<http://www.med.upenn.edu>

University of Pennsylvania, Philadelphia, Pennsylvania



**BOX 3. Hand-hygiene research agenda****Education and promotion**

- Provide health-care workers (HCWs) with better education regarding the types of patient care activities that can result in hand contamination and cross-transmission of microorganisms.
- Develop and implement promotion hand-hygiene programs in pregraduate courses.
- Study the impact of population-based education on hand-hygiene behavior.
- Design and conduct studies to determine if frequent glove use should be encouraged or discouraged.
- Determine evidence-based indications for hand cleansing (considering that it might be unrealistic to expect HCWs to clean their hands after every contact with the patient).
- Assess the key determinants of hand-hygiene behavior and promotion among the different populations of HCWs.
- Develop methods to obtain management support.
- Implement and evaluate the impact of the different components of multimodal programs to promote hand hygiene.

**Hand-hygiene agents and hand care**

- Determine the most suitable formulations for hand-hygiene products.
- Determine if preparations with persistent antimicrobial activity reduce infection rates more effectively than do preparations whose activity is limited to an immediate effect.
- Study the systematic replacement of conventional handwashing by the use of hand disinfection.
- Develop devices to facilitate the use and optimal application of hand-hygiene agents.
- Develop hand-hygiene agents with low irritancy potential.
- Study the possible advantages and eventual interaction of hand-care lotions, creams, and other barriers to help minimize the potential irritation associated with hand-hygiene agents.

**Laboratory-based and epidemiologic research and development**

- Develop experimental models for the study of cross-contamination from patient to patient and from environment to patient.
- Develop new protocols for evaluating the in vivo efficacy of agents, considering in particular short application times and volumes that reflect actual use in health-care facilities.
- Monitor hand-hygiene adherence by using new devices or adequate surrogate markers, allowing frequent individual feedback on performance.
- Determine the percentage increase in hand-hygiene adherence required to achieve a predictable risk reduction in infection rates.
- Generate more definitive evidence for the impact on infection rates of improved adherence to recommended hand-hygiene practices.
- Provide cost-effectiveness evaluation of successful and unsuccessful promotion campaigns.

**Part II. Recommendations****Categories**

These recommendations are designed to improve hand-hygiene practices of HCWs and to reduce transmission of pathogenic microorganisms to patients and personnel in health-care settings. This guideline and its recommendations are not intended for use in food processing or food-service establishments, and are not meant to replace guidance provided by FDA's Model Food Code.

As in previous CDC/HICPAC guidelines, each recommendation is categorized on the basis of existing scientific data, theoretical rationale, applicability, and economic impact. The CDC/HICPAC system for categorizing recommendations is as follows:

*Category IA.* Strongly recommended for implementation and strongly supported by well-designed experimental, clinical, or epidemiologic studies.

*Category IB.* Strongly recommended for implementation and supported by certain experimental, clinical, or epidemiologic studies and a strong theoretical rationale.

*Category IC.* Required for implementation, as mandated by federal or state regulation or standard.

*Category II.* Suggested for implementation and supported by suggestive clinical or epidemiologic studies or a theoretical rationale.

*No recommendation.* Unresolved issue. Practices for which insufficient evidence or no consensus regarding efficacy exist.

## Recommendations

1. Indications for handwashing and hand antisepsis
  - A. When hands are visibly dirty or contaminated with proteinaceous material or are visibly soiled with blood or other body fluids, wash hands with either a non-antimicrobial soap and water or an antimicrobial soap and water (IA) (66).
  - B. If hands are not visibly soiled, use an alcohol-based hand rub for routinely decontaminating hands in all other clinical situations described in items 1C–J (IA) (74,93,166,169,283,294,312,398). Alternatively, wash hands with an antimicrobial soap and water in all clinical situations described in items 1C–J (IB) (69-71,74).
  - C. Decontaminate hands before having direct contact with patients (IB) (68,400).
  - D. Decontaminate hands before donning sterile gloves when inserting a central intravascular catheter (IB) (401,402).
  - E. Decontaminate hands before inserting indwelling urinary catheters, peripheral vascular catheters, or other invasive devices that do not require a surgical procedure (IB) (25,403).
  - F. Decontaminate hands after contact with a patient's intact skin (e.g., when taking a pulse or blood pressure, and lifting a patient) (IB) (25,45,48,68).
  - G. Decontaminate hands after contact with body fluids or excretions, mucous membranes, nonintact skin, and wound dressings if hands are not visibly soiled (IA) (400).
  - H. Decontaminate hands if moving from a contaminated-body site to a clean-body site during patient care (II) (25,53).
  - I. Decontaminate hands after contact with inanimate objects (including medical equipment) in the immediate vicinity of the patient (II) (46,53,54).
  - J. Decontaminate hands after removing gloves (IB) (50,58,321).
  - K. Before eating and after using a restroom, wash hands with a non-antimicrobial soap and water or with an antimicrobial soap and water (IB) (404-409).
  - L. Antimicrobial-impregnated wipes (i.e., towelettes) may be considered as an alternative to washing hands with non-antimicrobial soap and water. Because they are not as effective as alcohol-based hand rubs or washing hands with an antimicrobial soap and water for reducing bacterial counts on the hands of HCWs, they are not a substitute for using an alcohol-based hand rub or antimicrobial soap (IB) (160,161).
  - M. Wash hands with non-antimicrobial soap and water or with antimicrobial soap and water if exposure to *Bacillus anthracis* is suspected or proven. The physical action of washing and rinsing hands under such circumstances is recommended because alcohols, chlorhexidine, iodophors, and other antiseptic agents have poor activity against spores (II) (120,172,224,225).
  - N. No recommendation can be made regarding the routine use of nonalcohol-based hand rubs for hand hygiene in health-care settings. Unresolved issue.
2. Hand-hygiene technique
  - A. When decontaminating hands with an alcohol-based hand rub, apply product to palm of one hand and rub hands together, covering all surfaces of hands and fingers, until hands are dry (IB) (288,410). Follow the manufacturer's recommendations regarding the volume of product to use.
  - B. When washing hands with soap and water, wet hands first with water, apply an amount of product recommended by the manufacturer to hands, and rub hands together vigorously for at least 15 seconds, covering all surfaces of the hands and fingers. Rinse hands with water and dry thoroughly with a disposable towel. Use towel to turn off the faucet (IB) (90-92,94,411). Avoid using hot water, because repeated exposure to hot water may increase the risk of dermatitis (IB) (254,255).
  - C. Liquid, bar, leaflet or powdered forms of plain soap are acceptable when washing hands with a non-antimicrobial soap and water. When bar soap is used, soap racks that facilitate drainage and small bars of soap should be used (II) (412-415).
  - D. Multiple-use cloth towels of the hanging or roll type are not recommended for use in health-care settings (II) (137,300).
3. Surgical hand antisepsis
  - A. Remove rings, watches, and bracelets before beginning the surgical hand scrub (II) (375,378,416).
  - B. Remove debris from underneath fingernails using a nail cleaner under running water (II) (14,417).

- C. Surgical hand antisepsis using either an antimicrobial soap or an alcohol-based hand rub with persistent activity is recommended before donning sterile gloves when performing surgical procedures (IB) (115,159,232,234,237,418).
  - D. When performing surgical hand antisepsis using an antimicrobial soap, scrub hands and forearms for the length of time recommended by the manufacturer, usually 2–6 minutes. Long scrub times (e.g., 10 minutes) are not necessary (IB) (117,156,205,207,238-241).
  - E. When using an alcohol-based surgical hand-scrub product with persistent activity, follow the manufacturer's instructions. Before applying the alcohol solution, prewash hands and forearms with a non-antimicrobial soap and dry hands and forearms completely. After application of the alcohol-based product as recommended, allow hands and forearms to dry thoroughly before donning sterile gloves (IB) (159,237).
4. Selection of hand-hygiene agents
    - A. Provide personnel with efficacious hand-hygiene products that have low irritancy potential, particularly when these products are used multiple times per shift (IB) (90,92,98,166,249). This recommendation applies to products used for hand antisepsis before and after patient care in clinical areas and to products used for surgical hand antisepsis by surgical personnel.
    - B. To maximize acceptance of hand-hygiene products by HCWs, solicit input from these employees regarding the feel, fragrance, and skin tolerance of any products under consideration. The cost of hand-hygiene products should not be the primary factor influencing product selection (IB) (92,93,166,274,276-278).
    - C. When selecting non-antimicrobial soaps, antimicrobial soaps, or alcohol-based hand rubs, solicit information from manufacturers regarding any known interactions between products used to clean hands, skin care products, and the types of gloves used in the institution (II) (174,372).
    - D. Before making purchasing decisions, evaluate the dispenser systems of various product manufacturers or distributors to ensure that dispensers function adequately and deliver an appropriate volume of product (II) (286).
    - E. Do not add soap to a partially empty soap dispenser. This practice of "topping off" dispensers can lead to bacterial contamination of soap (IA) (187,419).
  5. Skin care
    - A. Provide HCWs with hand lotions or creams to minimize the occurrence of irritant contact dermatitis associated with hand antisepsis or handwashing (IA) (272,273).
    - B. Solicit information from manufacturers regarding any effects that hand lotions, creams, or alcohol-based hand antiseptics may have on the persistent effects of antimicrobial soaps being used in the institution (IB) (174,420,421).
  6. Other Aspects of Hand Hygiene
    - A. Do not wear artificial fingernails or extenders when having direct contact with patients at high risk (e.g., those in intensive-care units or operating rooms) (IA) (350–353).
    - B. Keep natural nails tips less than 1/4-inch long (II) (350).
    - C. Wear gloves when contact with blood or other potentially infectious materials, mucous membranes, and nonintact skin could occur (IC) (356).
    - D. Remove gloves after caring for a patient. Do not wear the same pair of gloves for the care of more than one patient, and do not wash gloves between uses with different patients (IB) (50,58,321,373).
    - E. Change gloves during patient care if moving from a contaminated body site to a clean body site (II) (50,51,58).
    - F. No recommendation can be made regarding wearing rings in health-care settings. Unresolved issue.
  7. Health-care worker educational and motivational programs
    - A. As part of an overall program to improve hand-hygiene practices of HCWs, educate personnel regarding the types of patient-care activities that can result in hand contamination and the advantages and disadvantages of various methods used to clean their hands (II) (74,292,295,299).
    - B. Monitor HCWs' adherence with recommended hand-hygiene practices and provide personnel with information regarding their performance (IA) (74,276,292,295,299,306,310).
    - C. Encourage patients and their families to remind HCWs to decontaminate their hands (II) (394,422).
  8. Administrative measures
    - A. Make improved hand-hygiene adherence an institutional priority and provide appropriate

- administrative support and financial resources (IB) (74,75).
- B. Implement a multidisciplinary program designed to improve adherence of health personnel to recommended hand-hygiene practices (IB) (74,75).
- C. As part of a multidisciplinary program to improve hand-hygiene adherence, provide HCWs with a readily accessible alcohol-based hand-rub product (IA) (74,166,283,294,312).
- D. To improve hand-hygiene adherence among personnel who work in areas in which high workloads and high intensity of patient care are anticipated, make an alcohol-based hand rub available at the entrance to the patient's room or at the bedside, in other convenient locations, and in individual pocket-sized containers to be carried by HCWs (IA) (11,74,166,283,284,312,318,423).
- E. Store supplies of alcohol-based hand rubs in cabinets or areas approved for flammable materials (IC).

### Part III. Performance Indicators

1. The following performance indicators are recommended for measuring improvements in HCWs' hand-hygiene adherence:
- A. Periodically monitor and record adherence as the number of hand-hygiene episodes performed by personnel/number of hand-hygiene opportunities, by ward or by service. Provide feedback to personnel regarding their performance.
- B. Monitor the volume of alcohol-based hand rub (or detergent used for handwashing or hand antisepsis) used per 1,000 patient-days.
- C. Monitor adherence to policies dealing with wearing of artificial nails.
- D. When outbreaks of infection occur, assess the adequacy of health-care worker hand hygiene.

### References

- Rotter M. Hand washing and hand disinfection [Chapter 87]. In: Mayhall CG, ed. Hospital epidemiology and infection control. 2<sup>nd</sup> ed. Philadelphia, PA: Lippincott Williams & Wilkins, 1999.
- Labarraque AG. Instructions and observations regarding the use of the chlorides of soda and lime. Porter J, ed. [French] New Haven, CT: Baldwin and Treadway, 1829.
- Semmelweis I. Etiology, concept, and prophylaxis of childbed fever. Carter KC, ed. 1<sup>st</sup> ed. Madison, WI: The University of Wisconsin Press, 1983.
- Coppage CM. Hand washing in patient care [Motion picture]. Washington, DC: US Public Health Service, 1961.
- Steere AC, Mallison GF. Handwashing practices for the prevention of nosocomial infections. *Ann Intern Med* 1975;83:683-90.
- Garner JS, Favero MS. CDC guideline for handwashing and hospital environmental control, 1985. *Infect Control* 1986;7:231-43.
- Larson E. Guideline for use of topical antimicrobial agents. *Am J Infect Control* 1988;16:253-66.
- Larson EL, APIC Guidelines Committee. APIC guideline for handwashing and hand antisepsis in health care settings. *Am J Infect Control* 1995;23:251-69.
- Hospital Infection Control Practices Advisory Committee (HICPAC). Recommendations for preventing the spread of vancomycin resistance. *Infect Control Hosp Epidemiol* 1995;16:105-13.
- Garner JS, Hospital Infection Control Practices Advisory Committee. Guideline for isolation precautions in hospitals. *Infect Control Hosp Epidemiol* 1996;17:53-80.
- Pittet D, Mourouga P, Perneger TV, Members of the Infection Control Program. Compliance with handwashing in a teaching hospital. *Ann Intern Med* 1999;130:126-30.
- Boyce JM. It is time for action: improving hand hygiene in hospitals. *Ann Intern Med* 1999;130:153-5.
- Selwyn S. Microbiology and ecology of human skin. *Practitioner* 1980;224:1059-62.
- Price PB. Bacteriology of normal skin: a new quantitative test applied to a study of the bacterial flora and the disinfectant action of mechanical cleansing. *J Infect Dis* 1938;63:301-18.
- Larson E. Effects of handwashing agent, handwashing frequency, and clinical area on hand flora. *Am J Infect Control* 1984;11:76-82.
- Maki D. Control of colonization and transmission of pathogenic bacteria in the hospital. *Ann Intern Med* 1978;89(Pt 2):777-80.
- Larson EL, Norton Hughes CA, Pyrak JD, Sparks SM, Cagatay EU, Bartkus JM. Changes in bacterial flora associated with skin damage on hands of health care personnel. *Am J Infect Control* 1998;26:513-21.
- Sprunt K, Redman W, Leidy G. Antibacterial effectiveness of routine hand washing. *Pediatrics* 1973;52:264-71.
- Food and Drug Administration. Tentative final monograph for healthcare antiseptic drug products; proposed rule. *Federal Register* 1994;59:31441-52.
- Lowbury EJJ. Gram-negative bacilli on the skin. *Br J Dermatol* 1969;81(suppl 1):55-61.
- Noble WC. Distribution of the Micrococccaceae. *Br J Dermatol* 1969;81(suppl 1):27-31.
- McBride ME, Duncan WC, Bodey GP, McBride CM. Microbial skin flora of selected cancer patients and hospital personnel. *J Clin Microbiol* 1976;3:14-20.
- Casewell MW. Role of hands in nosocomial gram-negative infection. In: Maibach HI, Aly R, eds. Skin microbiology: relevance to clinical infection. New York, NY: Springer-Verlag, 1981.
- Larson EL, McGinley KJ, Foglia AR, Talbot GH, Leyden JJ. Composition and antimicrobial resistance of skin flora in hospitalized and healthy adults. *J Clin Microbiol* 1986;23:604-8.
- Ehrenkranz NJ, Alfonso BC. Failure of bland soap handwash to prevent hand transfer of patient bacteria to urethral catheters. *Infect Control Hosp Epidemiol* 1991;12:654-62.
- Sanderson PJ, Weisler S. Recovery of coliforms from the hands of nurses and patients: activities leading to contamination. *J Hosp Infect* 1992;21:85-93.
- Coello R, Jiménez J, García M, et al. Prospective study of infection, colonization and carriage of methicillin-resistant *Staphylococcus aureus* in an outbreak affecting 990 patients. *Eur J Clin Microbiol Infect Dis* 1994;13:74-81.



28. Sanford MD, Widmer AF, Bale MJ, Jones RN, Wenzel RP. Efficient detection and long-term persistence of the carriage of methicillin-resistant *Staphylococcus aureus*. Clin Infect Dis 1994;19:1123–8.
29. Bertone SA, Fisher MC, Mortensen JE. Quantitative skin cultures at potential catheter sites in neonates. Infect Control Hosp Epidemiol 1994;15:315–8.
30. Bonten MJM, Hayden MK, Nathan C, VanVoorhis J, et al. Epidemiology of colonisation of patients and environment with vancomycin-resistant enterococci. Lancet 1996;348:1615–9.
31. Larson EL, Cronquist AB, Whittier S, Lai L, Lyle CT, Della Latta P. Differences in skin flora between inpatients and chronically ill patients. Heart Lung 2000;29:298–305.
32. Polakoff S, Richards IDG, Parker MT, Lidwell OM. Nasal and skin carriage of *Staphylococcus aureus* by patients undergoing surgical operation. J Hyg (Lond) 1967;65:559–66.
33. Leyden JJ, McGinley KJ, Nordstrom KM, Webster GF. Skin microflora. J Invest Dermatol 1987;88:65s–72s.
34. Tuazon CU, Perez A, Kishaba T, Sheagren JN. *Staphylococcus aureus* among insulin-injecting diabetic patients. JAMA 1975;231:1272.
35. Kaplowitz LG, Comstock JA, Landwehr DM, Dalton HP, Mayhall CG. Prospective study of microbial colonization of the nose and skin and infection of the vascular access site in hemodialysis patients. J Clin Microbiol 1988;26:1257–62.
36. Aly R, Maibach HI, Shinefield HR. Microbial flora of atopic dermatitis. Arch Dermatol 1977;113:780–2.
37. Kirmani N, Tuazon CU, Murray HW, Parrish AE, Sheagren JN. *Staphylococcus aureus* carriage rate of patients receiving long-term hemodialysis. Arch Intern Med 1978;138:1657–9.
38. Goldblum SE, Ulrich JA, Goldman RS, Reed WP. Nasal and cutaneous flora among hemodialysis patients and personnel: quantitative and qualitative characterization and patterns of staphylococcal carriage. Am J Kidney Dis 1982;11:281–6.
39. Boelaert JR, Van Landuyt HW, Gordts BZ, De Baere YA, Messer SA, Herwaldt LA. Nasal and cutaneous carriage of *Staphylococcus aureus* in hemodialysis patients: the effect of nasal mupirocin. Infect Control Hosp Epidemiol 1996;17:809–11.
40. Zimakoff J, Pedersen FB, Bergen L, et al. *Staphylococcus aureus* carriage and infections among patients in four haemo- and peritoneal-dialysis centres in Denmark. J Hosp Infect 1996;33:289–300.
41. Bibel DJ, Greenbert JH, Cook JL. *Staphylococcus aureus* and the microbial ecology of atopic dermatitis. Can J Microbiol 1997;23:1062–8.
42. Noble WC. Dispersal of skin microorganisms. Br J Dermatol 1975;93:477–85.
43. Walter CW, Kundsins RB, Shilkret MA, Day MM. Spread of staphylococci to the environment. Antibiotics Annual 1959:952–7.
44. Boyce JM, Opal SM, Chow JW, et al. Outbreak of multidrug-resistant *Enterococcus faecium* with transferable *vanB* class vancomycin resistance. J Clin Microbiol 1994;32:1148–53.
45. McFarland LV, Mulligan ME, Kwok RYY, Stamm WE. Nosocomial acquisition of *Clostridium difficile* infection. N Engl J Med 1989;320:204–10.
46. Samore MH, Venkataraman L, DeGirolami PC, Levin E, Arbeit RD, Karchmer AW. Clinical and molecular epidemiology of sporadic and clustered cases of nosocomial *Clostridium difficile* diarrhea. Am J Med 1996;100:32–40.
47. Lidwell OM, Towers AG, Ballard J, Gladstone B. Transfer of microorganisms between nurses and patients in a clean air environment. J Appl Bacteriol 1974;37:649–56.
48. Casewell M, Phillips I. Hands as route of transmission for *Klebsiella* species. Br Med J 1977;2:1315–7.
49. Hall CB, Douglas RG. Modes of transmission of respiratory syncytial virus. J Pediatr 1981;99:100–2.
50. Olsen RJ, Lynch P, Coyle MB, Cummings J, Bokete T, Stamm WE. Examination gloves as barriers to hand contamination in clinical practice. JAMA 1993;270:350–3.
51. Pittet D, Dharan S, Touveneau S, Sauvan V, Perneger TV. Bacterial contamination of the hands of hospital staff during routine patient care. Arch Intern Med 1999;159:821–6.
52. Fox MK, Langner SB, Wells RW. How good are hand washing practices? Am J Nursing 1974;74:1676–8.
53. Ojajärvi J. Effectiveness of hand washing and disinfection methods in removing transient bacteria after patient nursing. J Hyg (Lond) 1980;85:193–203.
54. Boyce JM, Potter-Bynoe G, Chenevert C, King T. Environmental contamination due to methicillin-resistant *Staphylococcus aureus*: possible infection control implications. Infect Control Hosp Epidemiol 1997;18:622–7.
55. Hayden MK, Blom, DW, Lyle, EA, et al. The risk of hand and glove contamination by healthcare workers (HCWs) after contact with a VRE (+) patient (pt) or the pts environment (env) [Abstract K-1334]. Presented at the 41<sup>st</sup> Interscience Conference on Antimicrobial Agents and Chemotherapy Chicago: American Society for Microbiology, 2001.
56. Scott E, Bloomfield SF. The survival and transfer of microbial contamination via cloths, hands and utensils. J Appl Bacteriol 1990;68:271–8.
57. Bauer TM, Ofner E, Just HM, Just H, Daschner FD. An epidemiological study assessing the relative importance of airborne and direct contact transmission of microorganisms in a medical intensive care unit. J Hosp Infect 1990;15:301–9.
58. Tenorio AR, Badri SM, Sahgal NB, et al. Effectiveness of gloves in the prevention of hand carriage of vancomycin-resistant *Enterococcus* species by health care workers after patient care. Clin Infect Dis 2001;32:826–9.
59. Daschner FD. How cost-effective is the present use of antiseptics? J Hosp Infect 1988;11 (suppl A):227–35.
60. Knittle MA, Eitzman DV, Baer H. Role of hand contamination of personnel in the epidemiology of gram-negative nosocomial infections. J Pediatr 1975;86:433–7.
61. Ayliffe GAJ, Babb JR, Davies JG, Lilly HA. Hand disinfection: a comparison of various agents in laboratory and ward studies. J Hosp Infect 1988;11:226–43.
62. Strausbaugh LJ, Sewell DL, Ward TT, Pfaller MA, Heitzman T, Tjoelker R. High frequency of yeast carriage on hands of hospital personnel. J Clin Microbiol 1994;32:2299–300.
63. Marples RR, Towers AG. A laboratory model for the investigation of contact transfer of micro-organisms. J Hyg (Lond) 1979;82:237–48.
64. Mackintosh CA, Hoffman PN. An extended model for transfer of micro-organisms via the hands: differences between organisms and the effect of alcohol disinfection. J Hyg (Lond) 1984;92:345–55.
65. Patrick DR, Findon G, Miller TE. Residual moisture determines the level of touch-contact-associated bacterial transfer following hand washing. Epidemiol Infect 1997;119:319–25.
66. Larson E. A causal link between handwashing and risk of infection? Examination of the evidence. Infect Control Hosp Epidemiol 1988;9:28–36.
67. Larson E. Skin hygiene and infection prevention: more of the same or different approaches? Clin Infect Dis 1999;29:1287–94.

68. Mortimer EA Jr, Lipsitz PJ, Wolinsky E, Gonzaga AJ, Rammelkamp CH Jr. Transmission of staphylococci between newborns. *Am J Dis Child* 1962;104:289–95.
69. Maki DG. The use of antiseptics for handwashing by medical personnel. *J Chemother* 1989;1(suppl 1):3–11.
70. Massanari RM, Hierholzer WJ Jr. A crossover comparison of antiseptic soaps on nosocomial infection rates in intensive care units. *Am J Infect Control* 1984;12:247–8.
71. Doebbeling BN, Stanley GL, Sheetz CT, et al. Comparative efficacy of alternative hand-washing agents in reducing nosocomial infections in intensive care units. *N Engl J Med* 1992;327:88–93.
72. Webster J, Faoagali JL, Cartwright D. Elimination of methicillin-resistant *Staphylococcus aureus* from a neonatal intensive care unit after hand washing with triclosan. *J Paediatr Child Health* 1994;30:59–64.
73. Zafar AB, Butler RC, Reese DJ, Gaydos LA, Mennonona PA. Use of 0.3% triclosan (Bacti-Stat\*) to eradicate an outbreak of methicillin-resistant *Staphylococcus aureus* in a neonatal nursery. *Am J Infect Control* 1995;23:200–8.
74. Pittet D, Hugonnet S, Harbarth S, Mourouga P, Sauvan V, Touveneau S. Effectiveness of a hospital-wide programme to improve compliance with hand hygiene. *Lancet* 2000;356:1307–12.
75. Larson EL, Early E, Cloonan P, Sugrue S, Parides M. An organizational climate intervention associated with increased handwashing and decreased nosocomial infections. *Behav Med* 2000;26:14–22.
76. Fridkin SK, Pear SM, Williamson TH, Galgiani JN, Jarvis WR. The role of understaffing in central venous catheter-associated bloodstream infections. *Infect Control Hosp Epidemiol* 1996;17:150–8.
77. Vicca AF. Nursing staff workload as a determinant of methicillin-resistant *Staphylococcus aureus* spread in an adult intensive therapy unit. *J Hosp Infect* 1999;43:109–13.
78. Harbarth S, Sudre P, Dharan S, Cadenas M, Pittet D. Outbreak of *Enterobacter cloacae* related to understaffing, overcrowding, and poor hygiene practices. *Infect Control Hosp Epidemiol* 1999;20:598–603.
79. European Committee for Standardization. Chemical disinfectants and antiseptics—hygienic handrub—test method and requirements (phase 2/step 2) [European standard EN 1500]. Brussels, Belgium: Central Secretariat; 1997.
80. Kramer A, Rudolph P, Kampf G, Pittet D. Limited efficacy of alcohol-based hand gels. *Lancet* 2002;359:1489–90.
81. Sattar SA, Abebe M, Bueti AJ, Jampani H, Newman J, Hua S. Activity of an alcohol-based hand gel against human adeno-, rhino-, and rotaviruses using the fingerpad method. *Infect Control Hosp Epidemiol* 2000;21:516–9.
82. Wolff MH, Schmitt J, Rahaus M, König A. Hepatitis A virus: a test method for virucidal activity. *J Hosp Infect* 2001;48(suppl A):S18–S22.
83. Steinmann J. Some principles of virucidal testing. *J Hosp Infect* 2001;48(suppl A):S15–S17.
84. Gould D, Ream E. Assessing nurses' hand decontamination performance. *Nursing Times* 1993;89:47–50.
85. Quraishi ZA, McGuckin M, Blais FX. Duration of handwashing in intensive care units: a descriptive study. *Am J Infect Control* 1984;11:83–7.
86. Lund S, Jackson J, Leggett J, Hales L, Dworkin R, Gilbert D. Reality of glove use and handwashing in a community hospital. *Am J Infect Control* 1994;22:352–7.
87. Meengs MR, Giles BK, Chisholm CD, Cordell WH, Nelson DR. Hand washing frequency in an emergency department. *Ann Emerg Med* 1994;23:1307–12.
88. Larson E, McGeer A, Quraishi ZA, et al. Effect of an automated sink on handwashing practices and attitudes in high-risk units. *Infect Control Hosp Epidemiol* 1991;12:422–8.
89. Broughall JM, Marshman C, Jackson B, Bird P. An automatic monitoring system for measuring handwashing frequency. *J Hosp Infect* 1984;5:447–53.
90. Ojajärvi J, Mäkelä P, Rantasalo I. Failure of hand disinfection with frequent hand washing: a need for prolonged field studies. *J Hyg (Lond)* 1977;79:107–19.
91. Larson EL, Eke PI, Wilder MP, Laughon BE. Quantity of soap as a variable in handwashing. *Infect Control* 1987;8:371–5.
92. Larson E, Leyden JJ, McGinley KJ, Grove GL, Talbot GH. Physiologic and microbiologic changes in skin related to frequent handwashing. *Infect Control* 1986;7:59–63.
93. Larson EL, Eke PI, Laughon BE. Efficacy of alcohol-based hand rinses under frequent-use conditions. *Antimicrob Agents Chemother* 1986;30:542–4.
94. Larson EL, Laughon BE. Comparison of four antiseptic products containing chlorhexidine gluconate. *Antimicrob Agents Chemother* 1987;31:1572–4.
95. Meers PD, Yeo GA. Shedding of bacteria and skin squames after handwashing. *J Hyg (Lond)* 1978;81:99–105.
96. Winnefeld M, Richard MA, Drancourt M, Grobb JJ. Skin tolerance and effectiveness of two hand decontamination procedures in everyday hospital use. *Br J Dermatol* 2000;143:546–50.
97. Maki DG, Zilz MA, Alvarado CJ. Evaluation of the antibacterial efficacy of four agents for handwashing. In: Nelson JC, Grassi C, eds. Current chemotherapy and infectious disease proceedings of the 11<sup>th</sup> International Congress on Chemotherapy and the 19<sup>th</sup> ICACC. Washington, DC: American Society for Microbiology, 1979.
98. Boyce JM, Kelliher S, Vallande N. Skin irritation and dryness associated with two hand-hygiene regimens: soap-and-water handwashing versus hand antiseptics with an alcoholic hand gel. *Infect Control Hosp Epidemiol* 2000;21:442–8.
99. Sartor C, Jacomo V, Duvivier C, Tissot-Dupont H, Sambuc R, Drancourt M. Nosocomial *Serratia marcescens* infections associated with extrinsic contamination of a liquid nonmedicated soap. *Infect Control Hosp Epidemiol* 2000;21:196–9.
100. Walter CW. Editorial: disinfection of hands. *Am J Surg* 1965;109:691–3.
101. Gravens DL, Butcher HR Jr, Ballinger WF, Dewar NE. Septisol antiseptic foam for hands of operating room personnel: an effective antibacterial agent. *Surgery* 1973;73:360–7.
102. Eitzen HE, Ritter MA, French MLV, Gioe TJ. A microbiological in-use comparison of surgical hand-washing agents. *J Bone Joint Surg Am* 1979;61–A:403–6.
103. Minakuchi K, Yamamoto Y, Matsunaga K, et al. The antiseptic effect of a quick drying rubbing type povidone-iodine alcoholic disinfectant solution. *Postgrad Med J* 1993;69(suppl 3):S23–S26.
104. Babb JR, Davies JG, Ayliffe GAJ. A test procedure for evaluating surgical hand disinfection. *J Hosp Infect* 1991;18(suppl B):41–9.
105. Bellamy K, Alcock R, Babb JR, Davies JG, Ayliffe GA. A test for the assessment of 'hygienic' hand disinfection using rotavirus. *J Hosp Infect* 1993;24:201–10.
106. Ayliffe GAJ, Babb JR, Quraishi AH. A test for 'hygienic' hand disinfection. *J Clin Pathol* 1978;31:923–8.
107. Lilly HA, Lowbury EJJ, Wilkins MD. Detergents compared with each other and with antiseptics as skin 'degerming' agents. *J Hyg (Lond)* 1979;82:89–93.

108. Ulrich JA. Clinical study comparing hibistat (0.5% chlorhexidine gluconate in 70% isopropyl alcohol) and betadine surgical scrub (7.5% povidone-iodine) for efficacy against experimental contamination of human skin. *Curr Ther Res* 1982;31:27–30.
109. Bartzokas CA, Gibson MF, Graham R, Pinder DC. A comparison of triclosan and chlorhexidine preparations with 60 per cent isopropyl alcohol for hygienic hand disinfection. *J Hosp Infect* 1983;4:245–55.
110. Rotter ML, Koller W, Wewalka G, Werner HP, Ayliffe GAJ, Babb JR. Evaluation of procedures for hygienic hand-disinfection: controlled parallel experiments on the Vienna test model. *J Hyg (Lond)* 1986;96:27–37.
111. Kjrln H, Andersen BM. Handwashing and disinfection of heavily contaminated hands—effective or ineffective? *J Hosp Infect* 1992;21:61–71.
112. Namura S, Nishijima S, Asada Y. An evaluation of the residual activity of antiseptic handrub lotions: an ‘in use’ setting study. *J Dermatol* 1994;21:481–5.
113. Jarvis JD, Wynne CD, Enwright L, Williams JD. Handwashing and antiseptic-containing soaps in hospital. *J Clin Path* 1979;32:732–7.
114. Pereira LJ, Lee GM, Wade KJ. An evaluation of five protocols for surgical handwashing in relation to skin condition and microbial counts. *J Hosp Infect* 1997;36:49–65.
115. Larson EL, Butz AM, Gullette DL, Laughon BA. Alcohol for surgical scrubbing? *Infect Control Hosp Epidemiol* 1990;11:139–43.
116. Aly R, Maibach HI. Comparative study on the antimicrobial effect of 0.5% chlorhexidine gluconate and 70% isopropyl alcohol on the normal flora of hands. *Appl Environ Microbiol* 1979;37:610–3.
117. Galle PC, Homesley HD, Rhyne AL. Reassessment of the surgical scrub. *Surg Gynecol Obstet* 1978;147:215–8.
118. Rosenberg A, Alatary SD, Peterson AF. Safety and efficacy of the antiseptic chlorhexidine gluconate. *Surg Gynecol Obstet* 1976;143:789–92.
119. Ayliffe GAJ, Babb JR, Bridges K, et al. Comparison of two methods for assessing the removal of total organisms and pathogens from the skin. *J Hyg (Lond)* 1975;75:259–74.
120. Larson EL, Morton HE. Alcohols [Chapter 11]. In: Block SS, ed. *Disinfection, sterilization and preservation*. 4<sup>th</sup> ed. Philadelphia, PA: Lea and Febiger, 1991:642–54.
121. Price PB. Ethyl alcohol as a germicide. *Arch Surg* 1939;38:528–42.
122. Harrington C, Walker H. The germicidal action of alcohol. *Boston Medical and Surgical Journal* 1903;148:548–52.
123. Price PB. New studies in surgical bacteriology and surgical technic. *JAMA* 1938;111:1993–6.
124. Coulthard CE, Sykes G. The germicidal effect of alcohol with special reference to its action on bacterial spores. *Pharmaceutical Journal* 1936;137:79–81.
125. Pohle WD, Stuart LS. The germicidal action of cleaning agents—a study of a modification of Price’s procedure. *J Infect Dis* 1940;67:275–81.
126. Gardner AD. Rapid disinfection of clean unwashed skin: further experiments. *Lancet* 1948;760–3.
127. Sakuragi T, Yanagisawa K, Dan K. Bactericidal activity of skin disinfectants on methicillin-resistant *Staphylococcus aureus*. *Anesth Analg* 1995;81:555–8.
128. Kampf G, Jarosch R, Rüden H. Limited effectiveness of chlorhexidine based hand disinfectants against methicillin-resistant *Staphylococcus aureus* (MRSA). *J Hosp Infect* 1998;38:297–303.
129. Kampf G, Höfer M, Wendt C. Efficacy of hand disinfectants against vancomycin-resistant enterococci in vitro. *J Hosp Infect* 1999;42:143–50.
130. Platt J, Bucknall RA. The disinfection of respiratory syncytial virus by isopropanol and a chlorhexidine-detergent handwash. *J Hosp Infect* 1985;6:89–94.
131. Krilov LR, Harkness SH. Inactivation of respiratory syncytial virus by detergents and disinfectants. *Pediatr Infect Dis* 1993;12:582–4.
132. Sattar SA, Tetro J, Springthorpe VS, Giulivi A. Preventing the spread of hepatitis B and C viruses: where are germicides relevant? *Am J Infect Control* 2001;29:187–97.
133. Woolwine JD, Gerberding JL. Effect of testing method on apparent activities of antiviral disinfectants and antiseptics. *Antimicrob Agents Chemother* 1995;39:921–3.
134. Pillsbury DM, Livingood CS, Nichols AC. Bacterial flora of the normal skin: report on the effect of various ointment and solutions, with comments on the clinical significance of this study. *Arch Dermatol* 1942;45:61–80.
135. Lowbury EJJ, Lilly HA, Ayliffe GAJ. Preoperative disinfection of surgeons’ hands: use of alcoholic solutions and effects of gloves on skin flora. *Br Med J* 1974;4:369–72.
136. Lilly HA, Lowbury EJJ, Wilkins MD, Zaggy A. Delayed antimicrobial effects of skin disinfection by alcohol. *J Hyg (Lond)* 1979;82:497–500.
137. Ansari SA, Springthorpe VS, Sattar SA, Tostowaryk W, Wells GA. Comparison of cloth, paper, and warm air drying in eliminating viruses and bacteria from washed hands. *Am J Infect Control* 1991;19:243–9.
138. Ansari SA, Sattar SA, Springthorpe VS, Wells GA, Tostowaryk W. In vivo protocol for testing efficacy of hand-washing agents against viruses and bacteria: experiments with rotavirus and *Escherichia coli*. *Appl Environ Microbiol* 1989;55:3113–8.
139. Steinmann J, Nehr Korn R, Meyer A, Becker K. Two in-vivo protocols for testing virucidal efficacy of handwashing and hand disinfection. *Zentralbl Hyg Umweltmed.* 1995;196:425–36.
140. Mbithi JN, Springthorpe VS, Sattar SA. Comparative in vivo efficiencies of hand-washing agents against hepatitis A virus (HM-175) and poliovirus type 1 (Sabin). *Appl Environ Microbiol* 1993;59:3463–9.
141. Schurmann W, Eggers HJ. Antiviral activity of an alcoholic hand disinfectant: comparison of the in vitro suspension test with in vivo experiments on hands, and on individual fingertips. *Antiviral Res* 1983;3:25–41.
142. Larson E, Bobo L. Effective hand degerming in the presence of blood. *J Emerg Med* 1992;10:7–11.
143. Dineen P, Hildick-Smith G. Antiseptic care of the hands [Chapter 21]. In: Maibach HI, Hildick-Smith G, eds. *Skin bacteria and their role in infection*. New York: McGraw-Hill, 1965.
144. Lilly HA, Lowbury EJJ. Transient skin flora: their removal by cleansing or disinfection in relation to their mode of deposition. *J Clin Path* 1978;31:919–22.
145. Rotter M, Koller W, Wewalka G. Povidone-iodine and chlorhexidine gluconate-containing detergents for disinfection of hands. *J Hosp Infect* 1980;1:149–58.
146. Rotter ML. Hygienic hand disinfection. *Infect Control* 1984;1:18–22.
147. Blech MF, Hartemann P, Paquin JL. Activity of non antiseptic soaps and ethanol for hand disinfection. *Zentralbl Bakteriell Hyg [B]* 1985;181:496–512.

148. Leyden JJ, McGinley KJ, Kaminer MS, et al. Computerized image analysis of full-hand touch plates: a method for quantification of surface bacteria on hands and the effect of antimicrobial agents. *J Hosp Infect* 1991;18(suppl B):13–22.
149. Rotter ML, Koller W. Test models for hygienic handrub and hygienic handwash: the effects of two different contamination and sampling techniques. *J Hosp Infect* 1992;20:163–71.
150. Zaragoza M, Sallés M, Gomez J, Bayas JM, Trilla A. Handwashing with soap or alcoholic solutions? A randomized clinical trial of its effectiveness. *Am J Infect Control* 1999;27:258–61.
151. Paulson DS, Fendler EJ, Dolan MJ, Williams RA. A close look at alcohol gel as an antimicrobial sanitizing agent. *Am J Infect Control* 1999;27:332–8.
152. Cardoso CL, Pereira HH, Zequin JC, Guilhermetti M. Effectiveness of hand-cleansing agents for removing *Acinetobacter baumannii* strain from contaminated hands. *Am J Infect Control* 1999;27:327–31.
153. Casewell MW, Law MM, Desai N. A laboratory model for testing agents for hygienic hand disinfection: handwashing and chlorhexidine for the removal of klebsiella. *J Hosp Infect* 1988;12:163–75.
154. Wade JJ, Desai N, Casewell MW. Hygienic hand disinfection for the removal of epidemic vancomycin-resistant *Enterococcus faecium* and gentamicin-resistant *Enterobacter cloacae*. *J Hosp Infect* 1991;18:211–8.
155. Huang Y, Oie S, Kamiya A. Comparative effectiveness of hand-cleansing agents for removing methicillin-resistant *Staphylococcus aureus* from experimentally contaminated fingertips. *Am J Infect Control* 1994;22:224–7.
156. Lowbury EJJ, Lilly HA. Disinfection of the hands of surgeons and nurses. *Br Med J* 1960;1:5184.
157. Berman RE, Knight RA. Evaluation of hand antisepsis. *Arch Environ Health* 1969;18:781–3.
158. Rotter ML, Simpson RA, Koller W. Surgical hand disinfection with alcohols at various concentrations: parallel experiments using the new proposed European standards method. *Infect Control Hosp Epidemiol* 1998;19:778–81.
159. Hobson DW, Woller W, Anderson L, Guthery E. Development and evaluation of a new alcohol-based surgical hand scrub formulation with persistent antimicrobial characteristics and brushless application. *Am J Infect Control* 1998;26:507–12.
160. Jones MV, Rowe GB, Jackson B, Pritchard NJ. The use of alcoholic paper wipes for routine hand cleansing: results of trials in two hospitals. *J Hosp Infect* 1986;8:268–74.
161. Butz AM, Laughon BE, Gullette DL, Larson EL. Alcohol-impregnated wipes as an alternative in hand hygiene. *Am J Infect Control* 1990;18:70–6.
162. Ojajarvi J. Handwashing in Finland. *J Hosp Infect* 1991;18(suppl B):35–40.
163. Newman JL, Seitz JC. Intermittent use of an antimicrobial hand gel for reducing soap-induced irritation of health care personnel. *Am J Infect Control* 1990;18:194–200.
164. Rotter ML, Koller W, Neumann R. The influence of cosmetic additives on the acceptability of alcohol-based hand disinfectants. *J Hosp Infect* 1991;18 (suppl B):57–63.
165. Larson EL, Aiello AE, Heilman JM, et al. Comparison of different regimens for surgical hand preparation. *AORN J* 2001;73:412–20.
166. Larson EL, Aiello AE, Bastyr J, et al. Assessment of two hand hygiene regimens for intensive care unit personnel. *Crit Care Med* 2001;29:944–51.
167. Ophaswongse S, Maibach HI. Alcohol dermatitis: allergic contact dermatitis and contact urticaria syndrome: a review. *Contact Dermatitis* 1994;30:1–6.
168. Rilliet A, Hunziker N, Brun R. Alcohol contact urticaria syndrome (immediate-type hypersensitivity): case report. *Dermatologica* 1980;161:361–4.
169. Widmer AF. Replace hand washing with use of a waterless alcohol hand rub? *Clin Infect Dis* 2000;31:136–43.
170. Bryant KA, Pearce J, Stover B. Flash fire associated with the use of alcohol-based antiseptic agent [Letter]. *Am J Infect Control* 2002;30:256–7.
171. Hsueh PR, Teng LJ, Yang PC, Pan HL, Ho SW, Luh KT. Nosocomial pseudoepidemic caused by *Bacillus cereus* traced to contaminated ethyl alcohol from a liquor factory. *J Clin Microbiol* 1999;37:2280–4.
172. Denton GW. Chlorhexidine [Chapter 16]. In: Block SS, ed. *Disinfection, sterilization and reservation*. 4<sup>th</sup> ed. Philadelphia, PA: Lea and Febiger, 1991.
173. Narang HK, Codd AA. Action of commonly used disinfectants against enteroviruses. *J Hosp Infect* 1983;4:209–12.
174. Walsh B, Blakemore PH, Drabu YJ. The effect of handcream on the antibacterial activity of chlorhexidine gluconate. *J Hosp Infect* 1987;9:30–3.
175. Lowbury EJJ, Lilly HA. Use of 4% chlorhexidine detergent solution (Hibiscrub) and other methods of skin disinfection. *Br Med J* 1973;1:510–5.
176. Paulson DS. Comparative evaluation of five surgical hand scrub preparations. *AORN J* 1994;60:246–56.
177. Stringeni L, Lapomarda V, Lisi P. Occupational hand dermatitis in hospital environments. *Contact Dermatitis* 1995;33:172–6.
178. Marrie TJ, Costerton JW. Prolonged survival of *Serratia marcescens* in chlorhexidine. *Appl Environ Microbiol* 1981;42:1093–102.
179. McAllister TA, Lucas CE, Mocan H, et al. *Serratia marcescens* outbreak in a paediatric oncology unit traced to contaminated chlorhexidine. *Scott Med J* 1989;34:525–8.
180. Vigeant P, Loo VG, Bertrand C, et al. An outbreak of *Serratia marcescens* infections related to contaminated chlorhexidine. *Infect Control Hosp Epidemiol* 1998;19:791–4.
181. Vu-Thien H, Darbord JC, Moissenet D, et al. Investigation of an outbreak of wound infections due to *Alcaligenes xylosoxidans* transmitted by chlorhexidine in a burns unit. *Eur J Clin Microbiol* 1998;17:724–6.
182. Larson E, Talbot GH. An approach for selection of health care personnel handwashing agents. *Infect Control* 1986;7:419–24.
183. Davies J, Babb JR, Ayliffe GAJ, Wilkins MD. Disinfection of the skin of the abdomen. *Br J Surg* 1978;65:855–8.
184. Larson E, Mayur K, Laughon BA. Influence of two handwashing frequencies on reduction in colonizing flora with three handwashing products used by health care personnel. *Am J Infect Control* 1988;17:83–8.
185. Soulsby ME, Barnett JB, Maddox S. Brief report: the antiseptic efficacy of chlorxylenol-containing vs. chlorhexidine gluconate-containing surgical scrub preparations. *Infect Control* 1986;7:223–6.
186. Aly R, Maibach HI. Comparative antibacterial efficacy of a 2-minute surgical scrub with chlorhexidine gluconate, povidone-iodine, and chloroxylenol sponge-brushes. *Am J Infect Control* 1988;16:173–7.
187. Archibald LK, Corl A, Shah B, et al. *Serratia marcescens* outbreak associated with extrinsic contamination of 1% chlorxylenol soap. *Infect Control Hosp Epidemiol* 1997;18:704–9.
188. Lowbury EJJ, Lilly HA, Bull JP. Disinfection of hands: removal of resident bacteria. *Br Med J* 1963;1:1251–6.



189. Kundsinn RB, Walter CW. The surgical scrub—practical consideration. *Arch Surg* 1973;107:75–7.
190. Lockhart J. How toxic is hexachlorophene? *Pediatrics* 1972;50:229–35.
191. Shuman RM, Leech RW, Alvord EC Jr. Neurotoxicity of hexachlorophene in humans: II. a clinicopathological study of 46 premature infants. *Arch Neurol* 1975;32:320–5.
192. Dixon RE, Kaslow RA, Mallison GF, Bennett JV. Staphylococcal disease outbreaks in hospital nurseries in the United States—December 1971 through March 1972. *Pediatrics* 1973;51:413–6.
193. Kaslow RA, Dixon RE, Martin SM, et al. Staphylococcal disease related to hospital nursery bathing practices—a nationwide epidemiologic investigation. *Pediatrics* 1973;51:418–29.
194. American Academy of Pediatrics, American College of Obstetricians and Gynecologists. Guidelines for perinatal care. 4<sup>th</sup> ed. Elk Grove Village, IL: American Academy of Pediatrics; Washington, DC: American Academy of Obstetricians and Gynecologists, 1997.
195. Gottardi W. Iodine and iodine compounds [Chapter 8]. In: Block SS, ed. *Disinfection, sterilization and preservation*. 4<sup>th</sup> ed. Philadelphia, PA: Lea & Febiger; 1991.
196. Anderson RL. Iodophor antiseptics: intrinsic microbial contamination with resistant bacteria. *Infect Control Hosp Epidemiol* 1989;10:443–6.
197. Goldenheim PD. In vitro efficacy of povidone-iodine solution and cream against methicillin-resistant *Staphylococcus aureus*. *Postgrad Med J* 1993;69(suppl 3):S62–S65.
198. Traoré O, Fayard SF, Laveran H. An in-vitro evaluation of the activity of povidone-iodine against nosocomial bacterial strains. *J Hosp Infect* 1996;34:217–22.
199. McLure AR, Gordon J. In-vitro evaluation of povidone-iodine and chlorhexidine against methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect* 1992;21:291–9.
200. Davies JG, Babb JR, Bradley CR, Ayliffe GAJ. Preliminary study of test methods to assess the virucidal activity of skin disinfectants using poliovirus and bacteriophages. *J Hosp Infect* 1993;25:125–31.
201. Rotter ML. Chapter 79/Hand washing and hand disinfection. In: Mayhall CG, ed. *Hospital Epidemiology and Infection Control*. Baltimore, MD: Williams & Wilkins, 1996.
202. Wade JJ, Casewell MW. The evaluation of residual antimicrobial activity on hands and its clinical relevance. *J Hosp Infect* 1991;18 (suppl B):23–8.
203. Aly R, Maibach HI. Comparative evaluation of chlorhexidine gluconate (Hibiclens<sup>®</sup>) and povidone-iodine (E-Z Scrub<sup>®</sup>) sponge/brushes for presurgical hand scrubbing. *Curr Ther Res* 1983;34:740–5.
204. Herruzo-Cabrera R, Vizcaino-Alcaide MJ, Fdez-AciZero MJ. Usefulness of an alcohol solution of N-duopropenide for the surgical antisepsis of the hands compared with handwashing with iodine-povidone and chlorhexidine: clinical essay. *J Surgical Research* 2000;94:6–12.
205. Hingst V, Juditzki I, Heeg P, Sonntag HG. Evaluation of the efficacy of surgical hand disinfection following a reduced application time of 3 instead of 5 min. *J Hosp Infect* 1992;20:79–86.
206. Faoagali J, Fong J, George N, Mahoney P, O'Rourke V. Comparison of the immediate, residual, and cumulative antibacterial effects of Novaderm R, Novascrub R, Betadine Surgical Scrub, Hibiclens, and liquid soap. *Am J Infect Control* 1995;23:337–43.
207. Pereira LJ, Lee GM, Wade KJ. The effect of surgical handwashing routines on the microbial counts of operating room nurses. *Am J Infect Control* 1990;18:354–64.
208. Peterson AF, Rosenberg A. Comparative evaluation of surgical scrub preparations. *Surg Gynecol Obstet* 1978;146:63–5.
209. Berkelman RL, Holland BW, Anderson RL. Increased bactericidal activity of dilute preparations of povidone-iodine solutions. *J Clin Microbiol* 1982;15:635–9.
210. Merianos JJ. Quaternary ammonium antimicrobial compounds [Chapter 13]. In: Block SS, ed. *Disinfection, Sterilization, and Preservation*. 4<sup>th</sup> ed. Philadelphia, PA: Lea and Febiger; 1991.
211. Dixon RE, Kaslow RA, Mackel DC, Fulkerson CC, Mallison GF. Aqueous quaternary ammonium antiseptics and disinfectants: use and misuse. *JAMA* 1976;236:2415–7.
212. Sautter RL, Mattman LH, Legaspi RC. *Serratia marcescens* meningitis associated with a contaminated benzalkonium chloride solution. *Infect Control* 1984;5:223–5.
213. Oie S, Kamiya A. Microbial contamination of antiseptics and disinfectants. *Am J Infect Control* 1996;24:389–95.
214. Hayes RA, Trick WE, Vernon MO, et al. Comparison of three hand hygiene (HH) methods in a surgical intensive care unit (SICU) [Abstract K-1337]. Presented at the 41<sup>st</sup> Interscience Conference on Antimicrobial Agents and Chemotherapy Chicago, IL: American Society for Microbiology, 2001.
215. Dyer DL, Gerenraich KB, Wadhams PS. Testing a new alcohol-free hand sanitizer to combat infection. *AORN J* 1998;68:239–51.
216. Jones RD, Jampani HB, Newman JL, Lee AS. Triclosan: a review of effectiveness and safety in health care settings. *Am J Infect Control* 2000;28:184–96.
217. Ward WH, Holdgate GA, Rowsell S, et al. Kinetic and structural characteristics of the inhibition of enoyl (acyl carrier protein) reductase by triclosan. *Biochemistry* 1999;38:12514–25.
218. Heath RJ, Li J, Roland GE. Inhibition of the *Staphylococcus aureus* NADPH-dependent enoyl-acyl carrier protein reductase by triclosan and hexachlorophene. *J Biol Chem* 2000;275:4654–9.
219. Faoagali JL, George N, Fong J, Davy J, Dowser M. Comparison of the antibacterial efficacy of 4% chlorhexidine gluconate and 1% triclosan handwash products in an acute clinical ward. *Am J Infect Control* 1999;27:320–6.
220. Barry MA, Craven DE, Goularte TA, Lichtenberg DA. *Serratia marcescens* contamination of antiseptic soap containing triclosan: implications for nosocomial infection. *Infect Control* 1984;5:427–30.
221. Lowbury EJL, Lilly HA, Bull JP. Disinfection of hands: removal of transient organisms. *Br Med J* 1964;2:230–3.
222. Rotter ML. Semmelweis' sesquicentennial: a little-noted anniversary of handwashing. *Current Opinion in Infectious Diseases* 1998;11:457–60.
223. Manivannan G, Brady MJ, Cahalan PT, et al. Immediate, persistent and residual antimicrobial efficiency of Surfaccine<sup>™</sup> hand sanitizer [Abstract]. *Infection Control Hosp Epidemiol* 2000;21:105.
224. Gershenfeld L. Povidone-iodine as a sporicide. *Am J Pharm* 1962;134:79–81.
225. Russell AD. Chemical sporicidal and sporostatic agents [Chapter 22]. In: Block SS, ed. *Disinfection, sterilization and preservation*. 4<sup>th</sup> ed. Philadelphia, PA: Lea and Febiger, 1991.
226. Johnson S, Gerding DN, Olson MM, et al. Prospective, controlled study of vinyl glove use to interrupt *Clostridium difficile* nosocomial transmission. *Am J Med* 1990;88:137–40.
227. Russell AD. Mechanisms of bacterial insusceptibility to biocides. *Am J Infect Control* 2001;29:259–61.
228. Cookson BD, Bolton MC, Platt JH. Chlorhexidine resistance in methicillin-resistant *Staphylococcus aureus* or just an elevated MIC? An in vitro and in vivo assessment. *Antimicrob Agents Chemother* 1991;35:1997–2002.

229. McMurry LM, Oethinger M, Levy SB. Overexpression of *marA*, *soxS*, or *acrAB* produces resistance to triclosan in laboratory and clinical strains of *Escherichia coli*. *FEMS Microbiol Lett* 1998;166:305–9.
230. Chuanchuen R, Beinlich K, Hoang TT, et al. Cross-resistance between triclosan and antibiotics in *Pseudomonas aeruginosa* is mediated by multidrug efflux pumps: exposure of a susceptible mutant strain to triclosan selects *rfxB* mutants overexpressing MexCD-OprJ. *Antimicrob Agents Chemother* 2001;45:428–32.
231. Gröschel DHM, Pruett TL. Surgical antisepsis [Chapter 36]. In: Block SS, ed. *Disinfection, sterilization and preservation*. 4<sup>th</sup> ed. Philadelphia, PA: Lea and Febiger, 1991.
232. Boyce JM, Potter-Bynoe G, Opal SM, Dziobek L, Medeiros AA. A common-source outbreak of *Staphylococcus epidermidis* infections among patients undergoing cardiac surgery. *J Infect Dis* 1990;161:493–9.
233. Dewar NE, Gravens DL. Effectiveness of seprisol antiseptic foam as a surgical scrub agent. *Appl Microbiol* 1973;26:544–9.
234. Grinbaum RS, de Mendonça JS, Cardo DM. An outbreak of handscrubbing-related surgical site infections in vascular surgical procedures. *Infect Control Hosp Epidemiol* 1995;16:198–202.
235. AORN Recommended Practices Committee. Recommended practices for surgical hand scrubs. In: Fogg D, Parker N, Shevlin D, eds. *Standards, Recommended Practices, and Guidelines*. Denver, CO: AORN, 2001.
236. Larson E, Anderson JK, Baxendale L, Bobo L. Effects of a protective foam on scrubbing and gloving. *Am J Infect Control* 1993;21:297–301.
237. Mulberry G, Snyder AT, Heilman J, Pyrek J, Stahl J. Evaluation of a waterless, scrubless chlorhexidine gluconate/ethanol surgical scrub for antimicrobial efficacy. *Am J Infect Control* 2001;29:377–82.
238. Dineen P. An evaluation of the duration of the surgical scrub. *Surg Gynecol Obstet* 1969;129:1181–4.
239. O'Farrell DA, Kenny G, O'Sullivan M, Nicholson P, Stephens M, Hone R. Evaluation of the optimal hand-scrub duration prior to total hip arthroplasty. *J Hosp Infect* 1994;26:93–8.
240. O'Shaughnessy M, O'Malley VP, Corbett G, Given HF. Optimum duration of surgical scrub-time [Short note]. *Br J Surg* 1991;78:685–6.
241. Wheelock SM, Lookinland S. Effect of surgical hand scrub time on subsequent bacterial growth. *AORN J* 1997;65:1087–98.
242. Deshmukh N, Kjellberg SI, Kramer JW. A comparison of 5-minute povidone-iodine scrub and 1-minute povidone-iodine scrub followed by alcohol foam. *Military Medicine* 1998;163:145–7.
243. Kikuchi-Numagami K, Saishu T, Fukaya M, Kanazawa E, Tagami H. Irritancy of scrubbing up for surgery with or without a brush. *Acta Derm Venereol* 1999;79:230–2.
244. Dineen P. The use of a polyurethane sponge in surgical scrubbing. *Surg Gynecol Obstet* 1966;123:595–8.
245. Bornside GH, Crowder VH Jr, Cohn I Jr. A bacteriological evaluation of surgical scrubbing with disposable iodophor-soap impregnated polyurethane scrub sponges. *Surgery* 1968;64:743–51.
246. McBride ME, Duncan WC, Knox JM. An evaluation of surgical scrub brushes. *Surg Gynecol Obstet* 1973;137:934–6.
247. Berman RE, Knight RA. Evaluation of hand antisepsis. *Arch Environ Health* 1969;18:781–3.
248. Loeb MB, Wilcox L, Smaill F, Walter S, Duff Z. A randomized trial of surgical scrubbing with a brush compared to antiseptic soap alone. *Am J Infect Control* 1997;25:11–5.
249. Larson E, Friedman C, Cohran J, Treston-Aurand J, Green S. Prevalence and correlates of skin damage on the hands of nurses. *Heart Lung* 1997;26:404–12.
250. Tupker RA. Detergents and cleansers [Chapter 7]. In: van der Valk PGM, Maibach HI, eds. *The Irritant Contact Dermatitis Syndrome*. New York, NY: CRC Press, 1996.
251. Wilhelm KP. Prevention of surfactant-induced irritant contact dermatitis. *Curr Probl Dermatol* 1996;25:78–85.
252. de Haan P, Meester HHM, Bruynzeel DP. Irritancy of alcohols [Chapter 6]. In: van der Valk PGM, Maibach HI, eds. *The Irritant Contact Dermatitis Syndrome*. New York, NY: CRC Press, 1996.
253. Lübke J, Ruffieux C, van Melle G, Perrenoud D. Irritancy of the skin disinfectant n-propanol. *Contact Dermatitis* 2001;45:226–31.
254. qhlenschlaeger J, Friberg J, Ramsing D, Agner T. Temperature dependency of skin susceptibility to water and detergents. *Acta Derm Venereol* 1996;76:274–6.
255. Emilson A, Lindberg M, Forslind B. The temperature effect of in vitro penetration of sodium lauryl sulfate and nickel chloride through human skin. *Acta Derm Venereol* 1993;73:203–7.
256. de Groot AC. Contact allergy to cosmetics: causative ingredients. *Contact Dermatitis* 1987;17:26–34.
257. Schnuch A, Uter W, Geier J, Frosch PJ, Rustemeyer T. Contact allergies in healthcare workers—results from the IVDK. *Acta Derm Venereol* 1998;78:358–63.
258. Rastogi SC, Heydorn S, Johansen JD, Basketter DA. Fragrance chemicals in domestic and occupational products. *Contact Dermatitis* 2001;45:221–5.
259. Uter W, Schnuch A, Geier J, Pfahlberg A, Gefeller O. Association between occupation and contact allergy to the fragrance mix: a multifactorial analysis of national surveillance data. *Occup Environ Med* 2001;58:392–8.
260. Perrenoud D, Bircher A, Hunziker T, et al. Frequency of sensitization to 13 common preservatives in Switzerland. *Contact Dermatitis* 1994;30:276–9.
261. Kiec-Swierczynska M, Krecisz B. Occupational skin diseases among the nurses in the region of Lodz. *Int J Occup Med Environ Health* 2000;13:179–84.
262. Garvey LH, Roed-Petersen J, and Husum B. Anaphylactic reactions in anaesthetised patients—four cases of chlorhexidine allergy. *Acta Anaesthesiol Scand* 2001;45:1290–4.
263. Pham, NH, Weiner JM, Reisner GS, and Baldo BA. Anaphylaxis to chlorhexidine. Case report. Implication of immunoglobulin E antibodies and identification of an allergenic determinant. *Clin Exp Allergy* 2000;30:1001–7.
264. Nishioka K, Seguchi T, Yasuno H, Yamamoto T, Tominaga K. The results of ingredient patch testing in contact dermatitis elicited by povidone-iodine preparations. *Contact Dermatitis* 2000;42:90–4.
265. Wong CSM, Beck MH. Allergic contact dermatitis from triclosan in antibacterial handwashes. *Contact Dermatitis* 2001;45:307.
266. Guin JD, Goodman J. Contact urticaria from benzyl alcohol presenting as intolerance to saline soaks. *Contact Dermatitis* 2001;45:182–3.
267. Podda M, Zollner T, Grundmann-Kollmann M, Kaufmann R, Boehncke WF. Allergic contact dermatitis from benzyl alcohol during topical antimycotic treatment. *Contact Dermatitis* 1999;41:302–3.
268. Yesudian PD, King CM. Allergic contact dermatitis from stearyl alcohol in Efudix<sup>®</sup> cream. *Contact Dermatitis* 2001;45:313–4.

269. Aust LB, Maibach HI. Incidence of human skin sensitization to isostearyl alcohol in two separate groups of panelists. *Contact Dermatitis* 1980;6:269–71.
270. Funk JO, Maibach HI. Propylene glycol dermatitis: re-evaluation of an old problem. *Contact Dermatitis* 1994;31:236–41.
271. Hannuksela M. Moisturizers in the prevention of contact dermatitis. *Curr Probl Dermatol* 1996;25:214–20.
272. Berndt U, Wigger-Alberti W, Gabard B, Elsner P. Efficacy of a barrier cream and its vehicle as protective measures against occupational irritant contact dermatitis. *Contact Dermatitis* 2000;42:77–80.
273. McCormick RD, Buchman TL, Maki DG. Double-blind, randomized trial of scheduled use of a novel barrier cream and an oil-containing lotion for protecting the hands of health care workers. *Am J Infect Control* 2000;28:302–10.
274. Larson E, Killien M. Factors influencing handwashing behavior of patient care personnel. *Am J Infect Control* 1982;10:93–9.
275. Zimakoff J, Kjelsberg AB, Larsen SO, Holstein B. A multicenter questionnaire investigation of attitudes toward hand hygiene, assessed by the staff in fifteen hospitals in Denmark and Norway. *Am J Infect Control* 1992;20:58–64.
276. Mayer JA, Dubbert PM, Miller M, Burkett PA, Chapman SW. Increasing handwashing in an intensive care unit. *Infect Control* 1986;7:259–62.
277. Ojajärvi J. The importance of soap selection for routine hand hygiene in hospital. *J Hyg (Lond)* 1981;86:275–83.
278. Scott D, Barnes A, Lister M, Arkell P. An evaluation of the user acceptability of chlorhexidine handwash formulations. *J Hosp Infect* 1991;18(suppl B):51–5.
279. Taylor LJ. An evaluation of handwashing techniques—2. *Nursing Times* 1978;74:108–10.
280. Preston GA, Larson EL, Stamm WE. The effect of private isolation rooms on patient care practices, colonization and infection in an intensive care unit. *Am J Med* 1981;70:641–5.
281. Kaplan LM, McGuckin M. Increasing handwashing compliance with more accessible sinks. *Infect Control* 1986;7:408–10.
282. Freeman, J. Prevention of nosocomial infections by location of sinks for hand washing adjacent to the bedside [Abstract 60]. In: Program and abstracts of the 33<sup>rd</sup> Interscience Conference on Antimicrobial Agents and Chemotherapy. Washington, DC: American Society for Microbiology, 1993:130.
283. Bischoff WE, Reynolds TM, Sessler CN, Edmond MB, Wenzel RP. Handwashing compliance by health care workers. The impact of introducing an accessible, alcohol-based hand antiseptic. *Arch Intern Med* 2000;160:1017–21.
284. Pittet D. Compliance with hand disinfection and its impact on hospital-acquired infections. *J Hosp Infect* 2001;48(suppl A):S40–S46.
285. Wurtz R, Moye G, Jovanovic B. Handwashing machines, handwashing compliance, and potential for cross-contamination. *Am J Infect Control* 1994;22:228–30.
286. Kohan C, Ligi C, Dumigan DG, Boyce JM. The importance of evaluating product dispensers when selecting alcohol-based handrubs. *Am J Infect Control* 2002 (in press).
287. Boyce JM. Antiseptic technology: access, affordability, and acceptance. *Emerg Infect Diseases* 2001;7:231–3.
288. Taylor LJ. An evaluation of handwashing techniques—1. *Nursing Times* 1978:54–5.
289. Albert RK, Condie F. Hand-washing patterns in medical intensive-care units. *N Engl J Med* 1981;304:1465–6.
290. Larson E. Compliance with isolation technique. *Am J Infect Control* 1983;11:221–5.
291. Donowitz LG. Handwashing technique in a pediatric intensive care unit. *Am J Dis Child* 1987;141:683–5.
292. Conly JM, Hill S, Ross J, Lertzman J, Loule TJ. Handwashing practices in an intensive care unit: the effects of an educational program and its relationship to infection rates. *Am J Infect Control* 1989;17:330–9.
293. DeCarvalho M, Lopes JMA, Pellitteri M. Frequency and duration of handwashing in a neonatal intensive care unit. *Pediatr Infect Dis J* 1989;8:179–80.
294. Graham M. Frequency and duration of handwashing in an intensive care unit. *Am J Infect Control* 1990;18:77–80.
295. Dubbert PM, Dolce J, Richter W, Miller M, Chapman SW. Increasing ICU staff handwashing: effects of education and group feedback. *Infect Control Hosp Epidemiol* 1990;11:191–3.
296. Simmons B, Bryant J, Neiman K, Spencer L, Arheart K. The role of handwashing in prevention of endemic intensive care unit infections. *Infect Control Hosp Epidemiol* 1990;11:589–94.
297. Pettinger A, Nettleman MD. Epidemiology of isolation precautions. *Infect Control Hosp Epidemiol* 1991;12:303–7.
298. Lohr JA, Ingram DL, Dudley SM, Lawton EL, Donowitz LG. Hand washing in pediatric ambulatory settings: an inconsistent practice. *Am J Dis Child* 1991;145:1198–9.
299. Raju TNK, Kobler C. Improving handwashing habits in the newborn nurseries. *Am J Med.Sci* 1991;302:355–8.
300. Larson EL, McGinley KJ, Foglia A, et al. Handwashing practices and resistance and density of bacterial hand flora on two pediatric units in Lima, Peru. *Am J Infect Control* 1992;20:65–72.
301. Zimakoff J, Stormark M, Larsen SO. Use of gloves and handwashing behaviour among health care workers in intensive care units. A multicentre investigation in four hospitals in Denmark and Norway. *J Hosp Infect* 1993;24:63–7.
302. Pelke S, Ching D, Easa D, Melish ME. Gowning does not affect colonization or infection rates in a neonatal intensive care unit. *Arch Pediatr Adolesc Med* 1994;148:1016–20.
303. Gould D. Nurses' hand decontamination practice: results of a local study. *J Hosp Infect* 1994;28:15–30.
304. Shay DK, Maloney SA, Montecalvo M, et al. Epidemiology and mortality risk of vancomycin-resistant enterococcal bloodstream infections. *J Infect Dis* 1995;172:993–1000.
305. Berg DE, Hershov RC, Ramirez CA. Control of nosocomial infections in an intensive care unit in Guatemala City. *Clin Infect Dis* 1995;21:588–93.
306. Tibballs J. Teaching hospital medical staff to handwash. *Med J Aust* 1996;164:395–8.
307. Slaughter S, Hayden MK, Nathan C, et al. A comparison of the effect of universal use of gloves and gowns with that of glove use alone on acquisition of vancomycin-resistant enterococci in a medical intensive care unit. *Ann Intern Med* 1996;125:448–56.
308. Dorsey ST, Cydulka RK, Emerman CL. Is handwashing teachable?: failure to improve handwashing behavior in an urban emergency department. *Acad Emerg Med* 1996;3:360–5.
309. Watanakunakorn C, Wang C, Hazy J. An observational study of hand washing and infection control practices by healthcare workers. *Infect Control Hosp Epidemiol* 1998;19:858–60.

310. Avila-Agüero ML, UmaZa MA, Jiménez AL, Faingezicht I, París MM. Handwashing practices in a tertiary-care, pediatric hospital and the effect on an educational program. *Clin Perform Qual Health Care* 1998;6:70-2.
311. Kirkland KB, Weinstein JM. Adverse effects of contact isolation. *Lancet* 1999;354:1177-8.
312. Maury E, Alzieu M, Baudel JL, et al. Availability of an alcohol solution can improve hand disinfection compliance in an intensive care unit. *Am J Respir Crit Care Med* 2000;162:324-7.
313. Muto CA, Siström MG, Farr BM. Hand hygiene rates unaffected by installation of dispensers of a rapidly acting hand antiseptic. *Am J Infect Control* 2000;28:273-6.
314. Jarvis WR. Handwashing—the Semmelweis lesson forgotten? *Lancet* 1994;344:1311-2.
315. Larson E, Kretzer EK. Compliance with handwashing and barrier precautions. *J Hosp Infect* 1995;30(suppl):88-106.
316. Sproat LJ, Inglis TJJ. A multicentre survey of hand hygiene practice in intensive care units. *J Hosp Infect* 1994;26:137-48.
317. Kretzer EK, Larson EL. Behavioral interventions to improve infection control practices. *Am J Infect Control* 1998;26:245-53.
318. Voss A, Widmer AF. No time for handwashing!? Handwashing versus alcoholic rub: can we afford 100% compliance? *Infect Control Hosp Epidemiol* 1997;18:205-8.
319. Larson E. Handwashing and skin physiologic and bacteriologic aspects. *Infect Control* 1985;6:14-23.
320. Thompson BL, Dwyer DM, Ussery XT, Denman S, Vacek P, Schwartz B. Handwashing and glove use in a long-term care facility. *Infect Control Hosp Epidemiol* 1997;18:97-103.
321. Doebbeling BN, Pfaller MA, Houston AK, Wenzel RP. Removal of nosocomial pathogens from the contaminated glove. *Ann Intern Med* 1988;109:394-8.
322. McLane C, Chenelly S, Sylwestrak ML, Kirchoff KT. A nursing practice problem: failure to observe aseptic technique. *Am J Infect Control* 1983;11:178-82.
323. Pittet D. Improving compliance with hand hygiene in hospitals. *Infect Control Hosp Epidemiol* 2000;21:381-6.
324. Teare L, Handwashing Liaison Group. Hand washing: a modest measure—with big effects. *Br Med J* 1999;318:686.
325. Teare EL, Cookson B, French GL, et al. UK handwashing initiative. *J Hosp Infect* 1999;43:1-3.
326. Larson EL, Bryan JL, Adler LM, Blane C. A multifaceted approach to changing handwashing behavior. *Am J Infect Control* 1997;25:3-10.
327. Weeks A. Why I don't wash my hands between each patient contact [Letter]. *Br Med J* 1999;319:518.
328. Webster J. Handwashing in a neonatal intensive care nursery: product acceptability and effectiveness of chlorhexidine gluconate 4% and triclosan 1%. *J Hosp Infect* 1992;21:137-41.
329. Kelen GD, Green GB, Hexter DA, et al. Substantial improvement in compliance with universal precautions in an emergency department following institution of policy. *Arch Intern Med* 1991;151:2051-6.
330. Lundberg GD. Changing physician behavior in ordering diagnostic tests [Editorial]. *JAMA* 1998;280:2036.
331. Phillips DF. "New look" reflects changing style of patient safety enhancement. *JAMA* 1999;281:217-9.
332. Harbarth S, Martin Y, Rohner P, Henry N, Auckenthaler R, Pittet D. Effect of delayed infection control measures on a hospital outbreak of methicillin-resistant *Staphylococcus aureus*. *J Hosp Infect* 2000;46:43-9.
333. Early E, Battle K, Cantwell E, English J, Lavin JE, Larson E. Effect of several interventions on the frequency of handwashing among elementary public school children. *Am J Infect Control* 1998;26:263-9.
334. Butz AM, Larson E, Fosarelli P, Yolken R. Occurrence of infectious symptoms in children in day care homes. *Am J Infect Control* 1990;18:347-53.
335. Kimel LS. Handwashing education can decrease illness absenteeism. *J Sch Nurs* 1996;12:14-6, 18.
336. Master D, Hess Longe S, Dickson H. Scheduled hand washing in an elementary school population. *Fam Med* 1997;29:336-9.
337. Roberts L, Smith W, Jorm L, Patel M, Douglas RM, McGilchrist C. Effect of infection control measures on the frequency of upper respiratory infection in child care: a randomized, controlled trial. *Pediatrics* 2000;105:738-42.
338. Roberts L, Jorm L, Patel M, Smith W, Douglas RM, McGilchrist C. Effect of infection control measures on the frequency of diarrheal episodes in child care: a randomized, controlled trial. *Pediatrics* 2000;105:743-6.
339. Khan MU. Interruption of shigellosis by handwashing. *Trans R Soc Trop Med Hyg* 1982;76:164-8.
340. Shahid NS, Greenough WB, Samadi AR, Huq MI, Rahman N. Hand washing with soap reduces diarrhoea and spread of bacterial pathogens in a Bangladesh village. *J Diarrhoeal Dis Res* 1996;14:85-9.
341. Stanton BE, Clemens JD. An educational intervention for altering water-sanitation behaviors to reduce childhood diarrhea in urban Bangladesh. *Am J Epidemiol* 1987;125:292-301.
342. McGinley KJ, Larson EL, Leyden JJ. Composition and density of microflora in the subungual space of the hand. *J Clin Microbiol* 1988;26:950-3.
343. Hedderwick SA, McNeil SA, Lyons MJ, Kauffman CA. Pathogenic organisms associated with artificial fingernails worn by healthcare workers. *Infect Control Hosp Epidemiol* 2000;21:505-9.
344. Baumgardner CA, Maragos CS, Larson EL. Effects of nail polish on microbial growth of fingernails: dispelling sacred cows. *AORN J* 1993;58:84-8.
345. Wynd CA, Samstag DE, Lapp AM. Bacterial carriage on the fingernails of OR nurses. *AORN J* 1994;60:796-805.
346. Gross A, Cutright DE, D'Allessandro SM. Effect of surgical scrub on microbial population under the fingernails. *Am J Surg* 1979;138:463-7.
347. Pottinger J, Burns S, Manske C. Bacterial carriage by artificial versus natural nails. *Am J Infect Control* 1989;17:340-4.
348. McNeil SA, Foster CL, Hedderwick SA, Kauffman CA. Effect of hand cleansing with antimicrobial soap or alcohol-based gel on microbial colonization of artificial fingernails worn by health care workers. *Clin Infect Dis* 2001;32:367-72.
349. Rubin DM. Prosthetic fingernails in the OR. *AORN J* 1988;47:944-5, 948.
350. Moolenaar RL, Crutcher M, San Joaquin VH, et al. A prolonged outbreak of *Pseudomonas aeruginosa* in a neonatal intensive care unit: did staff fingernails play a role in disease transmission? *Infect Control Hosp Epidemiol* 2000;21:80-5.
351. Passaro DJ, Waring L, Armstrong R, et al. Postoperative *Serratia marcescens* wound infections traced to an out-of-hospital source. *J Infect Dis* 1997;175:992-5.
352. Foca M, Jakob K, Whittier S, et al. Endemic *Pseudomonas aeruginosa* infection in a neonatal intensive care unit. *N Engl J Med* 2000;343:695-700.



353. Parry MF, Grant B, Yukna M, et al. *Candida* osteomyelitis and diskitis after spinal surgery: an outbreak that implicates artificial nail use. *Clin Infect Dis* 2001;32:352–7.
354. Garner JS, Simmons BP. Guideline for isolation precautions in hospitals. *Infect Control* 1983;4(suppl 4):245–325.
355. CDC. Recommendations for prevention of HIV transmission in health-care settings. *MMWR* 1987;36(suppl 2S):3S–18S.
356. Occupational Safety and Health Administration. 29 CFR Part 1910.1030. Occupational exposure to bloodborne pathogens: final rule. *Federal Register* 1991;29CFR Part 1910:1030.
357. Hartstein AI, Denny MA, Morthland VH, LeMonte AM, Pfaller MA. Control of methicillin-resistant *Staphylococcus aureus* in a hospital and an intensive care unit. *Infect Control Hosp Epidemiol* 1995;16:405–11.
358. Maki DG, McCormick RD, Zilz MA, Stolz SM, Alvarado CJ. An MRSA outbreak in a SICU during universal precautions: new epidemiology for nosocomial MRSA: downside for universal precautions [Abstract 473]. In: Program and abstracts of the 30<sup>th</sup> Interscience Conference on Antimicrobial Agents and Chemotherapy. Washington, DC: American Society for Microbiology, 1990.
359. Kotilainen HR, Brinker JP, Avato JL, Gantz NM. Latex and vinyl examination gloves: quality control procedures and implications for health care workers. *Arch Intern Med* 1989;149:2749–53.
360. Reingold AL, Kane MA, Hightower AW. Failure of gloves and other protective devices to prevent transmission of hepatitis B virus to oral surgeons. *JAMA* 1988;259:2558–60.
361. Korniewicz DM, Laughon BE, Butz A, Larson E. Integrity of vinyl and latex procedures gloves. *Nurs Res* 1989;38:144–6.
362. DeGroot-Kosolcharoen J, Jones JM. Permeability of latex and vinyl gloves to water and blood. *Am J Infect Control* 1989;17:196–201.
363. Korniewicz DM, Kirwin M, Cresci K, Markut C, Larson E. In-use comparison of latex gloves in two high-risk units: surgical intensive care and acquired immunodeficiency syndrome. *Heart Lung* 1992;21:81–4.
364. Korniewicz DM, Kirwin M, Cresci K, et al. Barrier protection with examination gloves: double versus single. *Am J Infect Control* 1994;22:12–5.
365. Siström MG, Muto CA, Neal J, Strain BA, Farr BM. Glove leakage rates as a function of latex content and brand: caveat emptor [Abstract 24]. In: Program and abstracts of the 10<sup>th</sup> Annual Meeting of Society of Healthcare Epidemiology of America, Orlando, Florida, 1998.
366. Flanagan H, Farr B. Continued evaluation of glove leakage rates at the University of Virginia. Presented at the 11<sup>th</sup> Annual Meeting of the Society for Healthcare Epidemiology of America, Toronto, Canada, April 1, 2001.
367. Korniewicz DM, Laughon BE, Cyr WH, Lytle CD, Larson E. Leakage of virus through used vinyl and latex examination gloves. *J Clin Microbiol* 1990;28:787–8.
368. Rego A, Roley L. In-use barrier integrity of gloves: latex and nitrile superior to vinyl. *Am J Infect Control* 1999;27:405–10.
369. Fisher MD, Reddy VR, Williams FM, Lin KY, Thacker JG, Edlich RF. Biomechanical performance of powder-free examination gloves. *J Emerg Med* 1999;17:1011–8.
370. Edlich RF, Suber F, Neal JG, Jackson EM, Williams FM. Integrity of powder-free examination gloves to bacteriophage penetration. *J Biomed Mater Res* 1999;48:755–8.
371. Murray CA, Burke FJT, McHugh S. An assessment of the incidence of punctures in latex and non-latex dental examination gloves in routine clinical practice. *Br Dental Journal* 2001;190:377–80.
372. Jones RD, Jampani H, Mulberry G, Rizer RL. Moisturizing alcohol hand gels for surgical hand preparation. *AORN J* 2000;71:584–99.
373. Patterson JE, Vecchio J, Pantelick EL, et al. Association of contaminated gloves with transmission of *Acinetobacter calcoaceticus* var. *antitratrus* in an intensive care unit. *Am J Med* 1991;91:479–83.
374. Lowbury EJJ. Aseptic methods in the operating suite. *Lancet* 1968;1:705–9.
375. Hoffman PN, Cooke EM, McCarville MR, Emmerson AM. Microorganisms isolated from skin under wedding rings worn by hospital staff. *Br Med J* 1985;290:206–7.
376. Jacobson G, Thiele JE, McCune JH, Farrell LD. Handwashing: ring-wearing and number of microorganisms. *Nurs Res* 1985;34:186–8.
377. Hayes RA, Trick WE, Vernon MO, et al. Ring use as a risk factor (RF) for hand colonization in a surgical intensive care unit (SICU) [Abstract K-1333]. In: Program and abstracts of the 41<sup>st</sup> Interscience Conference on Antimicrobial Agents and Chemotherapy. Washington, DC: American Society for Microbiology, 2001.
378. Salisbury DM, Hutfilz P, Treen LM, Bollin GE, Gautam S. The effect of rings on microbial load of health care workers' hands. *Am J Infect Control* 1997;25:24–7.
379. Spire B, Barré-Sinoussi F, Montagnier L, Chermann JC. Inactivation of lymphadenopathy associated virus by chemical disinfectants. *Lancet* 1984;2:899–901.
380. Martin LS, McDougal JS, Loskoski SL. Disinfection and inactivation of the human T lymphotropic virus type III/lymphadenopathy-associated virus. *J Infect Dis* 1985;152:400–3.
381. Resnick L, Veren K, Salahuddin SZ, Tondreau S, Markham PD. Stability and inactivation of HTLV-III/LAV under clinical and laboratory environments. *JAMA* 1986;255:1887–91.
382. van Bueren J, Larkin DP, Simpson RA. Inactivation of human immunodeficiency virus type 1 by alcohols. *J Hosp Infect* 1994;28:137–48.
383. Montefiori DC, Robinson WE Jr, Modliszewski A, Mitchell WM. Effective inactivation of human immunodeficiency virus with chlorhexidine antiseptics containing detergents and alcohol. *J Hosp Infect* 1990;15:279–82.
384. Wood A, Payne D. The action of three antiseptics/disinfectants against enveloped and non-enveloped viruses. *J Hosp Infect* 1998;38:283–95.
385. Harbison MA, Hammer SM. Inactivation of human immunodeficiency virus by Betadine products and chlorhexidine. *J Acquir Immune Defic Syndr* 1989;2:16–20.
386. Lavelle GC, Gubbe SL, Neveaux JL, Bowden BJ. Evaluation of an antimicrobial soap formula for virucidal efficacy in vitro against human immunodeficiency virus in a blood-virus mixture. *Antimicrob Agents Chemother* 1989;33:2034–6.
387. Bond WW, Favero MS, Petersen NJ, Ebert JW. Inactivation of hepatitis B virus by intermediate-to-high level disinfectant chemicals. *J Clin Microbiol* 1983;18:535–8.
388. Kobayashi H, Tsuzuki M, Koshimizu K, et al. Susceptibility of hepatitis B virus to disinfectants or heat. *J Clin Microbiol* 1984;20:214–6.
389. Kurtz JB. Virucidal effect of alcohols against echovirus 11 [Letter]. *Lancet* 1979;1:496–7.
390. Sattar SA, Raphael RA, Lochnan H, Springthorpe VS. Rotavirus inactivation by chemical disinfectants and antiseptics used in hospitals. *Can J Microbiol* 1983;29:1464–9.
391. Larson E, Silberger M, Jakob K, et al. Assessment of alternative hand hygiene regimens to improve skin health among neonatal intensive care unit nurses. *Heart Lung* 2000;29:136–42.

392. Gould D, Chamberlain A. The use of a ward-based educational teaching package to enhance nurses' compliance with infection control procedures. *J Clin Nursing* 1997;6:55-67.
393. Aspöck C, Koller W. A simple hand hygiene exercise. *Am J Infect Control* 1999;27:370-2.
394. McGuckin M, Waterman R, Porten L, et al. Patient education model for increasing handwashing compliance [Practice forum]. *Am J Infect Control* 1999;27:309-14.
395. Khatib M, Jamaledine G, Abdallah A, Ibrahim Y. Hand washing and use of gloves while managing patients receiving mechanical ventilation in the ICU. *Chest* 1999;116:172-5.
396. Haley RW, Bregman DA. The role of understaffing and overcrowding in recurrent outbreaks of staphylococcal infection in a neonatal special-care unit. *J Infect Dis* 1982;145:875-85.
397. Pittet D, Boyce JM. Hand hygiene and patient care: pursuing the Semmelweis legacy. *Lancet Infectious Diseases* 2001;April:9-20.
398. Boyce JM. Scientific basis for handwashing with alcohol and other waterless antiseptic agents. In: Rutala WA, ed. *Disinfection, sterilization and antiseptics: principles and practices in healthcare facilities*. Washington, DC: Association for Professionals in Infection Control and Epidemiology, Inc, 2001.
399. O'Boyle CA, Henly SJ, Duckett LJ. Nurses' motivation to wash their hands: a standardized measurement approach. *Applied Nursing Research* 2001;14:136-45.
400. Semmelweis IP. *Die aetiologie, der begriff und die prophylaxis des kindbettfiebers*. Pest, Wien und Leipzig: CA Hartleben's Verlags-Expedition 1861.
401. Eggmann P, Harbarth S, Constantin MN, Touveneau S, Chevrolet JC, Pittet D. Impact of a prevention strategy targeted at vascular-access care on incidence of infections acquired in intensive care. *Lancet* 2000;355:1864-8.
402. Bull DA, Neumayer LA, Hunter GC, et al. Improved sterile technique diminishes the incidence of positive line cultures in cardiovascular patients. *J Surgical Research* 1992;52:106-10.
403. Hirschmann H, Fux L, Podusel J, et al. The influence of hand hygiene prior to insertion of peripheral venous catheters on the frequency of complications. *J Hosp Infect* 2001;49:199-203.
404. Drusin LM, Sohmer M, Groshen SL, Spiritos MD, Senterfit LB, Christenson WN. Nosocomial hepatitis A infection in a paediatric intensive care unit. *Arch Dis Child* 1987;62:690-5.
405. Doebbeling BN, Li N, Wenzel RP. An outbreak of hepatitis A among health care workers: risk factors for transmission. *Am J Public Health* 1993;83:1679-84.
406. Standaert SM, Hutcheson RH, Schaffner W. Nosocomial transmission of *Salmonella gastroenteritis* to laundry workers in a nursing home. *Infect Control Hosp Epidemiol* 1994;15:22-6.
407. Rodriguez EM, Parrott C, Rolka H, Monroe SS, Dwyer DM. An outbreak of viral gastroenteritis in a nursing home: importance of excluding ill employees. *Infect Control Hosp Epidemiol* 1996;17:587-92.
408. Schaffner W, Lefkowitz LB Jr, Goodman JS, Koenig MG. Hospital outbreak of infections with group A streptococci traced to an asymptomatic anal carrier. *N Engl J Med* 1969;280:1224-5.
409. Viglionese A, Nottebart VF, Bodman HA, Platt R. Recurrent group A streptococcal carriage in a health care worker associated with widely separated nosocomial outbreaks. *Am J Med* 1991;91(suppl 3B):329S-33S.
410. Ojajarvi J. An evaluation of antiseptics used for hand disinfection in wards. *J Hyg (Lond)* 1976;76:75-82.
411. Mermel LA, Josephson SL, Dempsey J, Parenteau S, Perry C, Magill N. Outbreak of *Shigella sonnei* in a clinical microbiology laboratory. *J Clin Microbiol* 1997;35:3163-5.
412. McBride ME. Microbial flora of in-use soap products. *Appl Environ Microbiol* 1984;48:338-41.
413. Kabara JJ, Brady MB. Contamination of bar soaps under "in use" condition. *J Environ Pathol Toxicol Oncol* 1984;5:1-14.
414. Heinze JE, Yackovich F. Washing with contaminated bar soap is unlikely to transfer bacteria. *Epidem Inf* 1988;101:135-42.
415. Bannan EA, Judge LF. Bacteriological studies relating to handwashing: 1. the inability of soap bars to transmit bacteria. *Am J Public Health* 1965;55:915-21.
416. Field EA, McGowan P, Pearce PK, Martin MV. Rings and watches: should they be removed prior to operative dental procedures? *J Dent* 1996;24:65-9.
417. Lowbury EJJ, Lilly HA. Gloved hand as applicator of antiseptic to operation sites. *Lancet* 1975;2:153-6.
418. AORN Recommended Practices Committee. Recommended practices for surgical hand scrubs. *AORN J* 1999;69:842-50.
419. Grohskopf LA, Roth VR, Feikin DR, et al. *Serratia liquefaciens* bloodstream infections from contamination of epoetin alfa at a hemodialysis center. *N Engl J Med* 2001;344:1491-7.
420. Dharan S, Hugonnet S, Sax H, Pittet D. Evaluation of interference of a hand care cream with alcohol-based hand disinfection. *Occup Environ Dermatol* 2001;49:81-4.
421. Heeg P. Does hand care ruin hand disinfection? *J Hosp Infect* 2001;48(suppl A):S37-S39.
422. McGuckin M, Waterman R, Storr J, et al. Evaluation of a patient-empowering hand hygiene programme in the U.K. *J Hosp Infect* 2001;48:222-7.
423. Girou E, Oppein F. Handwashing compliance in a French university hospital: new perspective with the introduction of hand-rubbing with a waterless alcohol-based solution. *J Hosp Infect* 2001;48(suppl A):S55-S57.

## Appendix

### Antimicrobial Spectrum and Characteristics of Hand-Hygiene Antiseptic Agents\*

| Group                             | Gram-positive bacteria | Gram-negative bacteria | Mycobacteria | Fungi | Viruses | Speed of action | Comments   |
|-----------------------------------|------------------------|------------------------|--------------|-------|---------|-----------------|--|
| Alcohols                          | +++                    | +++                    | +++          | +++   | +++     | Fast            | Optimum concentration 60%–95%; no persistent activity      |
| Chlorhexidine (2% and 4% aqueous) | +++                    | ++                     | +            | +     | +++     | Intermediate    | Persistent activity; rare allergic reactions               |
| Iodine compounds                  | +++                    | +++                    | +++          | ++    | +++     | Intermediate    | Causes skin burns; usually too irritating for hand hygiene |
| Iodophors                         | +++                    | +++                    | +            | ++    | ++      | Intermediate    | Less irritating than iodine; acceptance varies             |
| Phenol derivatives                | +++                    | +                      | +            | +     | +       | Intermediate    | Activity neutralized by nonionic surfactants               |
| Tricolsan                         | +++                    | ++                     | +            | —     | +++     | Intermediate    | Acceptability on hands varies                              |
| Quaternary ammonium compounds     | +                      | ++                     | —            | —     | +       | Slow            | Used only in combination with alcohols; ecologic concerns  |

**Note:** +++ = excellent; ++ = good, but does not include the entire bacterial spectrum; + = fair; — = no activity or not sufficient.

\*Hexachlorophene is not included because it is no longer an accepted ingredient of hand disinfectants.











# MMWR<sup>TM</sup>

## Morbidity and Mortality Weekly Report

Recommendations and Reports

October 25, 2002 / Vol. 51 / No. RR-16

### Continuing Education Activity Sponsored by CDC Guideline for Hand Hygiene in Health-Care Settings

#### Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force

#### EXPIRATION — October 25, 2004

You must complete and return the response form electronically or by mail by **October 25, 2004**, to receive continuing education credit. If you answer all of the questions, you will receive an award letter for 1.75 hours Continuing Medical Education (CME) credit; 0.15 Continuing Education Units (CEUs); 1.5 hours Certified Health Education Specialist (CHES) credit; or 1.9 contact

hours Continuing Nursing Education (CNE) credit. If you return the form electronically, you will receive educational credit immediately. If you mail the form, you will receive educational credit in approximately 30 days. No fees are charged for participating in this continuing education activity.

#### INSTRUCTIONS

##### By Internet

1. Read this *MMWR* (Vol. 51, RR-16), which contains the correct answers to the questions beginning on the next page.
2. Go to the *MMWR* Continuing Education Internet site at <http://www.cdc.gov/mmwr/cme/conted.html>.
3. Select which exam you want to take and select whether you want to register for CME, CEU, CHES, or CNE credit.
4. Fill out and submit the registration form.
5. Select exam questions. To receive continuing education credit, you must answer all of the questions. Questions with more than one correct answer will instruct you to "Indicate all that apply."
6. Submit your answers no later than **October 25, 2004**.
7. Immediately print your Certificate of Completion for your records.

##### By Mail or Fax

1. Read this *MMWR* (Vol. 50, RR-16), which contains the correct answers to the questions beginning on the next page.
2. Complete all registration information on the response form, including your name, mailing address, phone number, and e-mail address, if available.
3. Indicate whether you are registering for CME, CEU, CHES, or CNE credit.
4. Select your answers to the questions, and mark the corresponding letters on the response form. To receive continuing education credit, you must answer all of the questions. Questions with more than one correct answer will instruct you to "Indicate all that apply."
5. Sign and date the response form or a photocopy of the form and send no later than **October 25, 2004**, to  
Fax: 404-639-4198 Mail: MMWR CE Credit  
Office of Scientific and Health Communications  
Epidemiology Program Office, MS C-08  
Centers for Disease Control and Prevention  
1600 Clifton Rd, N.E.  
Atlanta, GA 30333
6. Your Certificate of Completion will be mailed to you within 30 days.

#### ACCREDITATION

**Continuing Medical Education (CME).** CDC is accredited by the Accreditation Council for Continuing Medical Education (ACCME) to provide continuing medical education for physicians. CDC designates this educational activity for a maximum of 1.75 hours in category 1 credit toward the AMA Physician's Recognition Award. Each physician should claim only those hours of credit that he/she actually spent in the educational activity.

**Continuing Education Unit (CEU).** CDC has been approved as an authorized provider of continuing education and training programs by the International Association for Continuing Education and Training and awards 0.15 Continuing Education Units (CEUs).

**Certified Health Education Specialist (CHES).** CDC is a designated provider of continuing education contact hours in health education by the National Commission for Health Education Credentialing, Inc. This program is a designated event for CHES to receive 1.5 hours in category 1 credit in health education. CDC provider number GA0082.

**Continuing Nursing Education (CNE).** This activity for 1.9 contact hours is provided by CDC, which is accredited as a provider of continuing education in nursing by the American Nurses Credentialing Center's Commission on Accreditation.

CENTERS FOR DISEASE CONTROL AND PREVENTION

SAFER • HEALTHIER • PEOPLE<sup>TM</sup>

### Goal and Objectives

This *MMWR* provides evidence-based recommendations for hand hygiene in health-care settings. These recommendations were developed by the Healthcare Infection Control Practices Advisory Committee (HICPAC), the Society for Healthcare Epidemiology of America, the Association for Professionals in Infection Control and Epidemiology, and the Infectious Diseases Society of America Hand Hygiene Task Force. The goal of this report is to provide guidance for clinicians and other health-care practitioners regarding strategies to improve hand-hygiene practices and reduce transmission of microorganisms in health-care settings. Upon completion of this educational activity, the reader should be able to 1) describe the indications for hand hygiene in health-care settings; 2) list the advantages of alcohol-based hand rubs; and 3) describe the barriers to hand hygiene in health-care settings.

**To receive continuing education credit, please answer all of the following questions.**

1. **Hand hygiene refers to . . .**
  - A. handwashing using plain soap and water.
  - B. using an antiseptic hand rub (e.g alcohol, chlorhexidine, iodine).
  - C. handwashing using antimicrobial soap and water.
  - D. all of the above.
2. **Hand hygiene adherence in health-care facilities might be improved by . . .**
  - A. providing personnel with individual containers of alcohol-based hand rubs.
  - B. providing personnel with hand lotions or creams.
  - C. providing personnel with feedback regarding hand-hygiene adherence/performance.
  - D. all of the above.
3. **Alcohol-based hand rubs have good or excellent antimicrobial activity against all of the following except . . .**
  - A. viruses.
  - B. fungi.
  - C. mycobacteria.
  - D. bacterial spores.
  - E. gram-positive and gram-negative bacteria.
4. **Alcohol-based hand rubs are indicated for all of the following clinical situations except . . .**
  - A. when the hands are visibly soiled.
  - B. preoperative cleaning of hands by surgical personnel.
  - C. before inserting urinary catheters, intravascular catheters, or other invasive devices.
  - D. after removing gloves.
5. **Each of the following statements regarding alcohol-based hand rubs is true except . . .**
  - A. alcohol-based hand rubs reduce bacterial counts on the hands of health-care personnel more effectively than plain soaps.
  - B. alcohol-based hand rubs can be made more accessible than sinks or other handwashing facilities.
  - C. alcohol-based hand rubs require less time to use than traditional handwashing.
  - D. alcohol-based hand rubs have been demonstrated to cause less skin irritation and dryness than handwashing using soap and water.
  - E. alcohol-based hand rubs are only effective if they are applied for  $\geq 60$  seconds.
6. **Which of the following statements regarding preoperative surgical hand antisepsis is true?**
  - A. Antimicrobial counts on hands are reduced as effectively with a 5-minute scrub as with a 10-minute scrub.
  - B. A brush or sponge must be used when applying the antiseptic agent to adequately reduce bacterial counts on hands.
  - C. Alcohol-based hand rubs for preoperative surgical scrub have been associated with increased surgical site infection rates.
  - D. A and B are true.
  - E. A and C are true.
7. **Antimicrobial-impregnated wipes (i.e., towelettes) . . .**
  - A. might be considered as an alternative to handwashing with plain soap and water.
  - B. are as effective as alcohol-based hands rubs.
  - C. are as effective as washing hands with antimicrobial soap and water.
  - D. A and C.
8. **The following statements regarding hand hygiene in health-care settings are true except . . .**
  - A. Overall adherence among health-care personnel is approximately 40%.
  - B. Poor adherence to hand-hygiene practice is a primary contributor to health-care-associated infection and transmission of antimicrobial-resistant pathogens.
  - C. Personnel wearing artificial nails or extenders have been linked to nosocomial outbreaks.
  - D. Hand hygiene is not necessary if gloves are worn.
9. **Indicate your work setting.**
  - A. State/local health department.
  - B. Other public health setting.
  - C. Hospital clinic/private practice.
  - D. Managed care organization.
  - E. Academic institution.
  - F. Other.
10. **Which best describes your professional activities?**
  - A. Patient care — emergency/urgent care department.
  - B. Patient care — inpatient.
  - C. Patient care — primary-care clinic or office.
  - D. Laboratory/pharmacy.
  - E. Public health.
  - F. Other.
11. **I plan to use these recommendations as the basis for . . . (Indicate all that apply.)**
  - A. health education materials.
  - B. insurance reimbursement policies.
  - C. local practice guidelines.
  - D. public policy.
  - E. other.
12. **Each month, approximately how many patients do you examine?**
  - A. None.
  - B. 1–5.
  - C. 6–20.
  - D. 21–50.
  - E. 51–100.
  - F. >100.
13. **How much time did you spend reading this report and completing the exam?**
  - A. 1–1.5 hours.
  - B. More than 1.5 hours but fewer than 2 hours.
  - C. 2–2.5 hours.
  - D. More than 2.5 hours.



- 14. After reading this report, I am confident I can describe the guidance for clinicians and other health-care practitioners regarding strategies to improve hand-hygiene practices and reduce transmission of microorganisms in health-care settings.
  - A. Strongly agree.
  - B. Agree.
  - C. Neither agree nor disagree.
  - D. Disagree.
  - E. Strongly disagree.
- 15. After reading this report, I am confident I can describe the indications for hand hygiene in health-care settings.
  - A. Strongly agree.
  - B. Agree.
  - C. Neither agree nor disagree.
  - D. Disagree.
  - E. Strongly disagree.
- 16. After reading this report, I am confident I can list the advantages of alcohol-based hand rubs.
  - A. Strongly agree.
  - B. Agree.
  - C. Neither agree nor disagree.
  - D. Disagree.
  - E. Strongly disagree.

- 17. After reading this report, I am confident I can describe the barriers to hand hygiene in health-care settings.
  - A. Strongly agree.
  - B. Agree.
  - C. Neither agree nor disagree.
  - D. Disagree.
  - E. Strongly disagree.
- 18. The objectives are relevant to the goal of this report.
  - A. Strongly agree.
  - B. Agree.
  - C. Neither agree nor disagree.
  - D. Disagree.
  - E. Strongly disagree.
- 19. The tables and text boxes are useful.
  - A. Strongly agree.
  - B. Agree.
  - C. Neither agree nor disagree.
  - D. Disagree.
  - E. Strongly disagree.
- 20. Overall, the presentation of the report enhanced my ability to understand the material.
  - A. Strongly agree.
  - B. Agree.
  - C. Neither agree nor disagree.
  - D. Disagree.
  - E. Strongly disagree.

**MMWR Response Form for Continuing Education Credit**  
**October 25, 2002/Vol. 51/No. RR-16**  
**Guideline for Hand Hygiene in Health-Care Settings**

**Recommendations of the Healthcare Infection Control Practices Advisory Committee and the HICPAC/SHEA/APIC/IDSA Hand Hygiene Task Force**

To receive continuing education credit, you must

1. provide your contact information;
2. indicate your choice of CME, CEU, CHES, or CNE credit;
3. answer all of the test questions;
4. sign and date this form or a photocopy;
5. submit your answer form by October 25, 2004.

Failure to complete these items can result in a delay or rejection of your application for continuing education credit.

*Detach or photocopy.*

- Check One*
- CME Credit
  - CEU Credit
  - CHES Credit
  - CNE Credit

Last Name \_\_\_\_\_ First Name \_\_\_\_\_

Street Address or P.O. Box \_\_\_\_\_

Apartment \_\_\_\_\_ or \_\_\_\_\_ Suite \_\_\_\_\_

City \_\_\_\_\_ State \_\_\_\_\_ ZIP Code \_\_\_\_\_

Phone Number \_\_\_\_\_ Fax Number \_\_\_\_\_

E-Mail Address \_\_\_\_\_

Fill in the appropriate blocks to indicate your answers. Remember, you must answer all of the questions to receive continuing education credit!

|   |   |
|---|---|
| 1. [ ] A [ ] B [ ] C [ ] D              | 14. [ ] A [ ] B [ ] C [ ] D [ ] E       |
| 2. [ ] A [ ] B [ ] C [ ] D              | 15. [ ] A [ ] B [ ] C [ ] D [ ] E       |
| 3. [ ] A [ ] B [ ] C [ ] D [ ] E        | 16. [ ] A [ ] B [ ] C [ ] D [ ] E       |
| 4. [ ] A [ ] B [ ] C [ ] D              | 17. [ ] A [ ] B [ ] C [ ] D [ ] E       |
| 5. [ ] A [ ] B [ ] C [ ] D [ ] E        | 18. [ ] A [ ] B [ ] C [ ] D [ ] E       |
| 6. [ ] A [ ] B [ ] C [ ] D [ ] E        | 19. [ ] A [ ] B [ ] C [ ] D [ ] E       |
| 7. [ ] A [ ] B [ ] C [ ] D              | 20. [ ] A [ ] B [ ] C [ ] D [ ] E       |
| 8. [ ] A [ ] B [ ] C [ ] D              | 21. [ ] A [ ] B [ ] C [ ] D [ ] E       |
| 9. [ ] A [ ] B [ ] C [ ] D [ ] E [ ] F  | 22. [ ] A [ ] B [ ] C [ ] D [ ] E       |
| 10. [ ] A [ ] B [ ] C [ ] D [ ] E [ ] F | 23. [ ] A [ ] B [ ] C [ ] D [ ] E [ ] F |
| 11. [ ] A [ ] B [ ] C [ ] D [ ] E       |   |
| 12. [ ] A [ ] B [ ] C [ ] D [ ] E [ ] F |   |
| 13. [ ] A [ ] B [ ] C [ ] D             |   |

Signature \_\_\_\_\_ Date I Completed Exam \_\_\_\_\_

**21. These recommendations will affect my practice.**

- A. Strongly agree.
- B. Agree.
- C. Neither agree nor disagree.
- D. Disagree.
- E. Strongly disagree.

**22. The availability of continuing education credit influenced my decision to read this report.**

- A. Strongly agree.
- B. Agree.
- C. Neither agree nor disagree.
- D. Disagree.
- E. Strongly disagree.

**23. How did you learn about this continuing education activity?**

- A. Internet.
- B. Advertisement (e.g., fact sheet, *MMWR* cover, newsletter, or journal).
- C. Coworker/supervisor.
- D. Conference presentation.
- E. *MMWR* subscription.
- F. Other.

Correct answers for questions 1–8  
1. D; 2. D; 3. D; 4. A; 5. E; 6. A; 7. A; 8. D.

## Hospital Infection Control Practices Advisory Committee Members

**Chair:** Robert A. Weinstein, M.D., Cook County Hospital, Chicago, Illinois.

**Co-Chair:** Jane D. Siegel, M.D., University of Texas Southwestern Medical Center, Dallas, Texas.

**Executive Secretary:** Michele L. Pearson, M.D. CDC, Atlanta, Georgia.

**Members:** Raymond Y.W. Chinn, M.D., Sharp Memorial Hospital, San Diego, California; Alfred DeMaria, Jr, M.D., Massachusetts Department of Public Health, Jamaica Plain, Massachusetts; Elaine L. Larson, Ph.D., Columbia University School of Nursing, New York, New York; James T. Lee, M.D., Ph.D., University of Minnesota, Minneapolis, Minnesota; Ramon E. Moncada, M.D., Coronado Physician's Medical Center, Coronado, California; William A. Rutala, Ph.D., University of North Carolina, Chapel Hill, North Carolina; William E. Scheckler, M.D., University of Wisconsin, Madison, Wisconsin; and Marjorie A. Underwood, Mt. Diablo Medical Center, Concord, California.

## Hand Hygiene Task Force Members

**Chair:** John M. Boyce, M.D., Hospital of Saint Raphael, New Haven, Connecticut.

**Members:** Barry M. Farr, M.D., University of Virginia, Charlottesville, Virginia; William R. Jarvis, M.D., CDC, Atlanta, Georgia; Elaine L. Larson, Ph.D., Columbia School of Nursing, New York, New York; Maryanne McGuckin, DrScEd, University of Pennsylvania, Philadelphia, Pennsylvania; Carol O'Boyle, Ph.D., Minnesota Department of Health, Minneapolis, Minnesota; Didier Pittet, M.D., University of Geneva, Geneva, Switzerland; Jane D. Siegel, M.D., University of Texas Southwestern Medical Center, Dallas, Texas; Marjorie A. Underwood, Mt. Diablo Medical Center, Concord, California; Andreas F. Widmer, M.D., University Hospitals, Basel, Switzerland; Jolynn Zeller, Avera St. Lukes Hospital, Aberdeen, South Dakota; and Beth H. Stover, Kosair Children's Hospital, Louisville, Kentucky.

## Disclosure of Financial Interests and Relationships

**John Boyce:** Research and educational grants from Gojo Industries; honorarium from Procter and Gamble; and consultant to Bode Chemical.

**Barry Farr:** Research support from Gojo Industries.

**Elaine Larson:** Received products for testing in research studies from Procter and Gamble, 3M Corporation, and Steris.

**Carol O'Boyle:** Honorarium from 3M Corporation.

All *MMWR* references are available on the Internet at <http://www.cdc.gov/mmwr>. Use the search function to find specific articles.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.

The *Morbidity and Mortality Weekly Report (MMWR)* series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy each week, send an e-mail message to [listserv@listserv.cdc.gov](mailto:listserv@listserv.cdc.gov). The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's Internet server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/publications/mmwr>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone 888-232-3228.

All material in the *MMWR* series is in the public domain and may be used and reprinted without permission; however, citation of the source is appreciated.